

"Discrete" vacuum geometry as a tool for Dirac fundamental quantization of Minkowskian Higgs model.

L. D. Lantsman

Wissenschaftliche Gesellschaft bei Jüdische Gemeinde zu Rostock,
Augusten Strasse, 20,
18055, Rostock, Germany;
Tel. 049-0381-799-07-24,
llantsman@freenet.de

May 18, 2008

Abstract

We demonstrate that assuming the "discrete" vacuum geometry in the Minkowskian Higgs model with vacuum BPS monopole solutions can justify the Dirac fundamental quantization of that model.

The important constituent of this quantization is getting various rotary effects, including collective solid rotations inside the physical BPS monopole vacuum, and just assuming the "discrete" vacuum geometry seems to be that thing able to justify these rotary effects.

More precisely, assuming the "discrete" geometry for the appropriate vacuum manifold implies the presence of thread topological defects (side by side with point hedgehog topological defects and walls between different topological domains) inside this manifold in the shape of specific (rectilinear) threads: gauge and Higgs fields located in the spatial region intimately near the axis z of the chosen (rest) reference frame.

This serves as the source of collective solid rotations proceeding inside the BPS monopole vacuum suffered the Dirac fundamental quantization.

It will be argued that indeed the first-order phase transition occurs in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac.

This comes to the coexistence of two thermodynamic phases inside the appropriate BPS monopole vacuum.

There are the thermodynamic phases of collective solid rotations and superfluid potential motions.

PACS: 12.38.Aw, 14.80.Bn, 14.80.Hv, 67.40.-w, 67.40.Bz, 68.18.Jk, 74.20.De, 74.25.-q.

Keywords: Non-Abelian Theory, BPS Monopole, Minkowski Space, Topological Defects, Phase transitions, Superfluidity, Superconductivity.

Contents

1	Introduction.	3
2	Statement of problem.	7
3	How one can introduce "discrete" geometry for degeneration spaces in Minkowskian Higgs models.	20
4	How topologically nontrivial threads arise inside the discrete non-Abelian vacuum manifold.	32
5	Specific of Higgs phase in Dirac quantisation scheme.	49

1 Introduction.

In the recent studies [1, 2, 3], the Gauss-shell reduction of the Minkowskian Higgs model with vacuum BPS monopole solutions was discussed.

Such Gauss-shell reduction is, indeed, the particular case of the Dirac fundamental quantization [4] of gauge theories, when such (Minkowskian) gauge model is written down in terms of topological Dirac variables [5] A_i^D ($i = 1, 2$): transverse and gauge invariant (i.e. automatically physical) functionals of gauge fields.

As a result of this Gauss-shell reduction, the picture of the BPS monopole vacuum as a medium possessing manifest superfluid properties and various rotary effects was given.

More exactly, superfluid properties of the BPS monopole vacuum suffered the Dirac fundamental quantization [4] come to its potentiality, described correctly by the *Bogomol'nyi* [1, 2, 6, 7, 8, 9, 10, 11],

$$\mathbf{B} = \pm D\Phi, \quad (1.1)$$

and *Gribov ambiguity* [2, 7, 8, 12],

$$[D_i^2(\Phi_k^{(0)})]^{ab}\Phi_{(0)b} = 0, \quad (1.2)$$

equations.

These Eqs. implicate the vacuum "magnetic" field $\mathbf{B}$ and Higgs vacuum BPS monopole modes Φ (in particular, topologically trivial modes $\Phi_{(0)a}$).

$\Phi_k^{(0)}$ are Yang-Mills (YM) vacuum BPS monopole modes, and D is the (covariant) derivative.

The connection between the Bogomol'nyi and Gribov ambiguity equations is realised via the Bianchi identity $D B = 0$ (the latter identity is equivalent to say that the vacuum "magnetic" field $\mathbf{B}$ is "transverse").

In Ref. [6] it was pointed out to the transparent analogy between the BPS monopole vacuum and a liquid helium II specimen, possessing the manifest superfluidity.

This analogy comes, in particular, to the mathematical resemblance between the Bogomol'nyi equation (1.1) and the expression [6, 13]

$$\mathbf{v}_0 = \frac{\hbar}{m} \nabla \Phi(t, \mathbf{r}), \quad (1.3)$$

for the critical velocity $\mathbf{v}_0$ of superfluid potential motions inside a liquid helium II specimen ¹.

¹ $v_0 = \min (\epsilon/p)$ for the ratio of the energy ϵ and momentum p for quantum excitations spectrum in the given liquid helium II specimen. Herewith [14] at velocities of the liquid exceeding a *critical velocity* $v_0 = \min (\epsilon/p)$ for the ratio of the energy ϵ and momentum p for quantum excitations spectrum in the liquid helium II, the dissipation of the liquid helium energy occurs via arising excitation quanta with momenta $\mathbf{p}$ directed antiparallel to the velocity vector $\mathbf{v}$. Such dissipation of the liquid helium energy becomes advantageous [15] just at

$$\epsilon + \mathbf{p} \cdot \mathbf{v} < 0 \implies \epsilon - p v < 0.$$

Eq. (1.3) implicates the phase $\Phi(t, \mathbf{r})$ of the complex-value helium Bose condensate wave function $\Xi(t, \mathbf{r}) \in C$, given as [6, 13]

$$\Xi(t, \mathbf{r}) = \sqrt{n_0(t, \mathbf{r})} e^{i\Phi(t, \mathbf{r})}, \quad (1.4)$$

with $n_0(t, \mathbf{r})$ being the number of particles in the ground energy state $\epsilon = 0$.

m is the mass of a helium atom.

On the other hand, for any potential motion, the condition [6]

$$\text{rot grad } \Phi = 0 \quad (1.5)$$

is always fulfilled (to within a constant) for a scalar field Φ .

In particular, the potentiality condition (1.5) is fulfilled for the crucial velocity $\mathbf{v}_0$, (1.3), of superfluid potential motions inside a liquid helium II, i.e.

$$\text{rot } \mathbf{v}_0 = 0.$$

This implies the absence of vortices and friction forces inside the superfluid component in a liquid helium II specimen.

The similar conclusion may be drawn also about the vacuum "magnetic" field $\mathbf{B}$ in the Minkowskian Higgs model with vacuum BPS monopole solutions, described correctly by the Bogomol'nyi equation (1.1) [1, 2, 6, 7, 8, 9, 10, 11] (with the correction that in the latter case the operator grad would be replaced with the [covariant] derivative D).

Additionally, the Gribov ambiguity equation (Gribov equation) (1.2) the Minkowskian BPS monopole vacuum as an incompressible medium.

This fact was discussed in Ref. [2]; it comes to the presence of typical topological invariants in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac [4].

There are the magnetic charge $\mathbf{m}$ and the degree of the map referring to the $U(1) \subset SU(2)$ embedding.

At resolving the YM Gauss law constraint [1, 2, 3, 7, 8, 12]

$$\frac{\delta W}{\delta A_0^a} = 0 \iff [D^2(A)]^{ac} A_{0c} = D_i^{ac}(A) \partial_0 A_c^i \quad (1.6)$$

in terms of topological Dirac variables A_i^D , satisfying the gauge [5]

$$D_i^{ab}(\hat{A}^D) \partial_0(\hat{A}_i^D) \equiv 0, \quad (1.7)$$

the right-hand side of this constraint vanishes, and it becomes the homogeneous equation [3, 7, 8, 12]

$$[D_i^2(\Phi^{(0)})]^{ac} A_{0c} = 0, \quad (1.8)$$

permitting, in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac [4], the family of zero mode solutions [12, 16],

$$A_0^c(t, \mathbf{x}) = \dot{N}(t) \Phi_{(0)}^c(\mathbf{x}) \equiv Z^c, \quad (1.9)$$

implicating the topological variable $\dot{N}(t)$ and Higgs (topologically trivial) vacuum Higgs BPS monopole modes $\Phi_0^a(\mathbf{x})$.

The topological variables $\dot{N}(t)$ (respectively, $N(t)$), may be specified [3, 7, 8, 12] via the relation

$$\begin{aligned}\nu[A_0, \Phi^{(0)}] &= \frac{g^2}{16\pi^2} \int_{t_{\text{in}}}^{t_{\text{out}}} dt \int d^3x F_{\mu\nu}^a \tilde{F}^{a\mu\nu} = \frac{\alpha_s}{2\pi} \int d^3x F_{i0}^a B_i^a(\Phi^{(0)}) [N(t_{\text{out}}) - N(t_{\text{in}})] \\ &= N(t_{\text{out}}) - N(t_{\text{in}}) = \int_{t_{\text{in}}}^{t_{\text{out}}} dt \dot{N}(t),\end{aligned}\tag{1.10}$$

taking account of the natural duality between the tensors F_{i0}^a and F_{ij}^a .

Herewith $\nu[A_0, \Phi^{(0)}]$ is referred to as the vacuum Chern-Simons functional, implicating the asymptotical states "in" and "out" taking in the time instants t_{in} and t_{out} , respectively.

As it was discussed in Ref. [3] (repeating the arguments [17]), it may be set

$$t_{\text{in}} \rightarrow -\infty; \quad t_{\text{out}} \rightarrow \infty.$$

The nontrivial topological dynamics inherent in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac comes to the specific Josephson effect [17] in the enumerated model, i.e. to the existence of collective solid rotations inside the *physical* BPS monopole vacuum.

These collective solid rotations are described correctly by the free rotator action functional

$$W_N = \int d^4x \frac{1}{2} (F_{0i}^c)^2 = \int dt \frac{\dot{N}^2 I}{2}\tag{1.11}$$

involving the rotary momentum [5]

$$I = \int_V d^3x (D_i^{ac}(\Phi_k^0) \Phi_{0c})^2 = \frac{4\pi^2 \epsilon}{\alpha_s} = \frac{4\pi^2}{\alpha_s^2} \frac{1}{V \langle B^2 \rangle}.\tag{1.12}$$

The YM coupling constant

$$\alpha_s = \frac{g^2}{4\pi(\hbar c)^2}$$

enters this expression for I together with the typical size ϵ [7, 8] of BPS monopoles and the vacuum expectation value $\langle B^2 \rangle$ of the "magnetic" field.

In Ref. [3] there was argued that $\dot{N}(t)$ play the role of angular velocities of collective solid rotations inside the Minkowskian Higgs vacuum suffered the Dirac fundamental quantization [4].

Meanwhile, the action functional (1.11) contains vacuum "electric" monopoles (in the terminology [7, 8])

$$F_{i0}^a = \dot{N}(t) D_i^{ac}(\Phi_k^{(0)}) \Phi_{(0)c}(\mathbf{x})\tag{1.13}$$

and is associated with the real spectrum

$$P_N = \dot{N}I = 2\pi k + \theta; \quad \theta \in [-\pi, \pi]; \quad (1.14)$$

of the topological momentum P_N .

The one of manifestations of the Josephson effect [17] occurring inside the Minkowskian Higgs vacuum suffered the Dirac fundamental quantization are the existence of never vanishing (excluding the value $\theta = 0$ of the θ -angle) vacuum "electric" fields ("electric" monopoles) (1.13) in the shape [5]

$$F_{i0}^a \equiv E_i^a = \dot{N}(t) (D_i(\Phi_k^{(0)}) \Phi_{(0)})^a = P_N \frac{\alpha_s}{4\pi^2\epsilon} B_i^a(\Phi_{(0)}) = (2\pi k + \theta) \frac{\alpha_s}{4\pi^2\epsilon} B_i^a(\Phi_{(0)}). \quad (1.15)$$

Any vacuum "electric" field (1.15) achieves its minimum in the zero topological sector of the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac:

$$(E_i^a)_{\min} = \theta \frac{\alpha_s}{4\pi^2\epsilon} B_i^a; \quad -\pi \leq \theta \leq \pi. \quad (1.16)$$

Such is, briefly, the picture of the (topologically nontrivial) dynamics inherent in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac [4].

This picture seems to be correct at least at the absolute zero temperature $T = 0$, when collective solid rotations inside the BPS monopole vacuum proceed in the "non-stop" regime [17] and "friction forces" between this BPS monopole vacuum and its surroundings are absent.

In Ref. [3], there was asserted that "geometrically", collective solid rotations inside the BPS monopole vacuum suffered the Dirac fundamental quantization [4] are, indeed, rotations around the infinitely narrow cylinder of the effective diameter $\sim \epsilon$ (with ϵ being the typical size of BPS monopoles) along the axis z of the chosen (rest) reference frame ².

In the present study we attempt to ground this assertion.

This will be associated immediately with assuming the "discrete vacuum geometry" [1, 3]

$$R_{YM} \equiv SU(2)/U(1) \simeq \mathbf{Z} \otimes G_0/U_0,$$

with

$$\pi_1(U_0) = \pi_1(G_0) = 0$$

and

$$SU(2) \equiv G; \quad U(1) \equiv U,$$

for the degeneration space (vacuum manifold) in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac, involving [9] the existence of (rectilinear, topologically nontrivial) threads inside the mentioned vacuum manifold, just located intimately near the axis z of the chosen (rest) reference frame.

² For the first time, the idea of "cylinder topologies" in (non-Abelian) gauge theories was proposed, probably in the paper [18] (where also the topological dynamical variable $N(t)$ was introduced).

Vortices generated by such threads, similar to those [15] one can observe in a (rested) liquid helium II specimen, may be identified naturally with collective solid rotations inside the BPS monopole vacuum suffered the Dirac fundamental quantization [4].

Unlike the spatial region near the axis z , far off this axis (including the region $|\mathbf{x}| \rightarrow \infty$), superfluid potential motions inside the BPS monopole vacuum occur.

These motions are described correctly by the Bogomol'nyi, (1.1), and Gribov, (1.2), equations.

The coexistence of (delimited somewhat in the space) collective solid rotations and superfluid potential motions inside the BPS monopole vacuum suffered the Dirac fundamental quantization is thus on hand [3].

It is the sign of the first-order phase transition occurring in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac.

This phase transition is additional to the second-order one associated with the spontaneous breakdown of the initial $SU(2)$ gauge symmetry down to the $U(1)$ one.

In the present study we reveal the role of the discrete geometry assumed for the vacuum manifold R_{YM} in this first-order phase transition just as in the series of physical effects (including collective solid vacuum rotations) taking place in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac.

2 Statement of problem.

The ensuing exposition is organized as follows.

In *Section 3* (repeating partially the theses of Ref. [1]) we, utilizing the general theory of topological defects (see e.g. §Φ1 in [9]), construct explicitly the degeneration space (vacuum manifold) R_{YM} for the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac [4].

As we have seen already, it is possible to postulate the look [1] for this manifold, implicating the discrete multiplier $\mathbf{Z}$.

Such look of the YM vacuum manifold R_{YM} is destined by the discrete factorisation [1] of the initial, $SU(2)$, and residual, $U(1)$, gauge symmetries groups in the enumerated model:

$$G \equiv SU(2) \simeq G_0 \otimes \mathbf{Z}; \quad H \equiv U(1) \simeq U_0 \otimes \mathbf{Z}.$$

In *Section 2*, repeating the arguments [9], we discover three kinds of topological defects inside the vacuum manifold R_{YM} . There are thread and point hedgehog (stable) defects and domain walls between different topological sectors of R_{YM} .

Herewith thread topological defects are of a great importance for us as those determined physical rotary effects and the first-order phase transition taking place in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac. We shall make sure in this soon.

As it is well known [9], the sufficient condition for thread topological defects to exist in a gauge theory involving the spontaneous breakdown of the initial symmetry group is

the isomorphism

$$\pi_1(R) = \pi_0(H) \neq 0 \quad (2.1)$$

between the appropriate homotopical groups of the degeneration space R and residual symmetry group H in the considered gauge theory.

This isomorphism is correct, in particular, for the vacuum manifold R_{YM} .

We ground this utilizing the holonomies group arguments (developed in Ref. [7]; see also [9]).

The existence of point hedgehog topological defects inside the same degeneration space R is controlled by the isomorphism [6, 9]

$$\pi_2(R) = \pi_1(H) \neq 0. \quad (2.2)$$

These topological defects are always associated, in (Minkowskian) Higgs non-Abelian models, with various monopole solutions (typical of which will be us discussed below). Geometrically, they are located at the origin of coordinates.

Besides two said types of topological defects, the YM vacuum manifold R_{YM} possesses the some more kind of topological defects. There are domain walls between different topological sectors of this manifold.

The cause of said domain walls is in the manifest "discrete" geometry of the Minkowskian YM vacuum manifold R_{YM} [9]:

$$\pi_0(R_{YM}) = \mathbf{Z}.$$

Indeed, the width of a domain wall is roughly proportional to the inverse of the lowest mass of all the physical particles being present in the (gauge) model considered [19].

In the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac it is just the typical size $\epsilon \sim (m/\sqrt{\lambda})^{-1}$ of BPS monopoles.

Thus in the Bogomol'nyi-Prasad-Sommerfeld (BPS) limit [6, 7, 8, 9, 10]

$$m \rightarrow 0; \quad \lambda \rightarrow 0 \quad (2.3)$$

for the Higgs mass m and selfinteraction constant λ , widths of domain walls inside R_{YM} can take arbitrary values, including infinite ones.

Actually [5, 7, 8], ϵ is the function of the spatial volume $V \sim r^3$:

$$\epsilon(V) \sim V^{-1} \propto r^{-3}.$$

Thus walls between topological domains become in fact infinite (of the typical width $\epsilon(0) \rightarrow \infty$) in the spatial region $r = 0$ (including the infinite narrow cylinder of the effective diameter $\epsilon(\infty) \rightarrow 0$ with its axis symmetry z).

Vice verse, as $|\mathbf{x}| \rightarrow \infty$, domain walls become infinitely thin (of the typical width $\epsilon(\infty) \rightarrow 0$).

In Ref. [3], it was argued (implicating the general QFT arguments) that asymptotical (vacuum) states "in" and "out" (cf. (1.10)) in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac would be separated by the infinite time interval $T \rightarrow \infty$ [17].

Assuming infinite widths for domain walls inside the vacuum manifold R_{YM} in the spatial region intimately near the axis z of the chosen (rest) reference frame is in a good agreement with this setting $T \rightarrow \infty$.

Section 4 of the present study we devote to the discussion about (topologically non-trivial) rectilinear threads containing inevitably inside the vacuum manifold R_{YM} .

It will be shown, repeating the arguments [9], that there are YM fields

$$A_\theta(\rho, \theta, z) \equiv A_\mu \partial x^\mu / \partial \theta$$

These fields may be always represented in the z -invariant shape as [9]

$$A_\theta(\rho, \theta, z) = \exp(iM\theta) A_\theta(\rho) \exp(-iM\theta), \quad (2.4)$$

with M being the generator of the group G_1 of rigid rotations compensating changes in the vacuum (Higgs-YM) configuration (Φ^a, A_μ^a) at rotations around the axis z of the chosen (rest) reference frame.

In (2.4),

$$A_\theta(\rho) = M + \beta(\rho),$$

where the function $\beta(\rho)$ approaches zero as $\rho \rightarrow \infty$.

The elements of G_1 can be set as [9]

$$g_\theta = \exp(iM\theta). \quad (2.5)$$

YM fields A_θ are manifestly invariant with respect to shifts along the axis z .

It is important that rectilinear threads A_θ don't coincide with vacuum YM BPS monopole solutions, and, on the contrary, there are, indeed, gaps between directions of "magnetic" tensions vectors: $\mathbf{B}_1$,

$$|\mathbf{B}_1| \sim \partial_\rho A_\theta(\rho, \theta, z), \quad (2.6)$$

and $\mathbf{B}$, given by the Bogomol'nyi equation (1.1) and diverging as r^{-2} at the origin of coordinates [10].

These gaps just testify in favour the above discussed first-order phase transition occurring in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac [4].

In the Higgs sector of that model, there exist [9] also z -invariant (vacuum) Higgs solutions in a (small) neighbourhood of the origin of coordinates:

$$\Phi^{(n)}(\rho, \theta, z) = \exp(M\theta) \phi(\rho) \quad (n \in \mathbf{Z}), \quad \nabla_\mu \phi(\rho) \leq \text{const } \rho^{-1+\delta}; \quad \delta > 0; \quad (2.7)$$

that can join (in a smooth wise) vacuum Higgs BPS monopoles, belonging to the same topology n and disappearing [20] at the origin of coordinates.

Herewith, speaking "in a smooth wise", we imply that the covariant derivative $D\Phi$ of any vacuum Higgs field $\Phi_a^{(n)}$ merges with the covariant derivative of such a vacuum Higgs BPS monopole solution.

Following [9], the vacuum z -invariant, i.e. *axially symmetric*, (Higgs-YM) configuration (Φ^a, A_μ^a) (given via Eqs. (2.4) and (2.7), possessing, as it can be demonstrated [9], a finite linear energy density and obeying the appropriate equations of motions) can be treated as a rectilinear thread solution (called also *the rectilinear thread vortex* or *the rectilinear thread*).

The same terminology "rectilinear threads" is applicable (as we have seen this above with the example of Eq. (2.4) [9]) to z -invariant (axially symmetric) YM fields A_θ .

Obvious locating (2.4) of (topologically nontrivial) threads A_θ at the origin of coordinates (actually, in the spatial region $\rho \rightarrow 0$; the same is correctly also for Higgs thread modes Φ^a due to (2.7)) permits the natural (somewhat alternative) geometrical interpretation of (topologically nontrivial) threads as infinitely narrow tubes around the axis z over which the family of vortex solutions (Φ^a, A_μ^a) is specified actually (disappearing rapidly outside this spatial region).

The requirement for vacuum Higgs fields $\Phi_a^{(n)}$ to be smooth is quite natural if the goal is pursued, in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac [4], to justify various rotary effects inherent in this model.

In particular, vacuum "electric" monopoles (1.15) [5] are directly proportional to $D_i(\Phi_k^{(0)}) \Phi_{(0)}$.

These vacuum "electric" monopoles, in turn, enter explicitly the action functional (1.11), describing, in the Dirac fundamental quantization scheme [4], collective solid rotations inside the Minkowskian BPS monopole vacuum.

Such (smooth) sawing together appropriate vacuum Higgs modes $\Phi^{(n)}$ (which are [9] specific thread rectilinear vortices) and BPS monopoles is intended to remove the *seem-ing* contradiction between the manifest superfluid properties of the Minkowskian BPS monopole vacuum (suffered the Dirac fundamental quantization [4]), setting by the Bogomolny'i, (1.1), and Gribov ambiguity, (1.2), equations.

Moreover, one can assert [16] that

$$D B \sim D E = 0 \quad (2.8)$$

for vacuum "magnetic" and "electric" tensions in the quested Minkowskian Higgs model, i.e. that these tensions are, indeed, "transverse" vectors colinear each other.

This just implies the potential nature of the "electric" tension $\mathbf{E}$, that can be perceived as the above contradiction, on the face of it. Note that Eq. (1.15) [5] just reflects this colinearity.

Going out from this contradiction seems to be just in locating (topologically nontrivial) threads in the infinitely narrow cylinder of the effective diameter ϵ around the axis z and in joining (in a smooth wise) vacuum Higgs fields $\Phi_a^{(n)}$ and BPS monopole solutions.

In this case collective solid rotations (vortices) inside the Minkowskian BPS monopole vacuum, occurring actually in that spatial region around the axis z and described correctly by the action functional (1.11), become quite "legitimate", and simultaneously, the Gauss law constraint (1.8) is satisfied outward this region with smooth vacuum "electric" monopoles E_i^a [5], (1.15). Latter ones can be treated as thread solutions since vacuum

Higgs fields $\Phi_a^{(n)}$ are such (the same assertion is correct, of course, for Higgs vacuum BPS monopoles).

On the other hand, in the region of thread topological defects inside the discrete vacuum manifold R_{YM} , Eq. (2.8) is violated since the vacuum "magnetic" field $\mathbf{B}$ suffers a break in this region according to the arguments [10] (the vacuum "magnetic" field $\mathbf{B}$ set for BPS monopole solutions via the Bogomol'nyi equation (1.1) [9, 10, 11] diverges as r^{-2} at the origin of coordinates).

In other words, the plot of $D\Phi$ behaves as r^{-2} at the origin of coordinates for vacuum Higgs BPS monopole modes Φ (whence $DB \sim D^2\Phi = O(r^{-3})$ therein), while the plot of $D\Phi$ for Higgs thread solutions Φ [9] joins the plot for the transverse vacuum "electric" field $E \sim D\Phi$ (2.8) [16] (in the latter case Φ are vacuum BPS monopole solutions, specified via the Bogomol'nyi equation (1.1)). Such is the geometrical picture connected with Eq. (2.8).

The said is the one more argument in favour of the first-order phase transition occurring in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac and coming therein to the coexistence of two thermodynamic phases: the thermodynamic phases of collective solid rotations and superfluid potential motions, inside the appropriate Minkowskian BPS monopole vacuum.

Herewith the enough clear-cut picture can be observed how the enumerated thermodynamic phases are distributed inside the discrete vacuum manifold R_{YM} .

Thread topological defects (vortices), associated with rectilinear threads A_θ , are located intimately near the axis z of the chosen (rest) reference frame. Actually, they refer to the cylinder of the effective diameter ϵ with z serving its symmetry axis.

Simultaneously, superfluid potential motions refer to the spatial region out of this cylinder, including the spatial region $|\mathbf{x}| \rightarrow \infty$ (corresponding to the infrared region of the momentum space).

The important consequence of the presence of rectilinear threads A_θ in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac [4] and involving the "discrete" vacuum geometry for the vacuum manifold R_{YM} is the effect [9] of annihilating two equal magnetic charges $\mathbf{m}_1 = \mathbf{m}_2 = \mathbf{m}(n) \neq 0$ ($n \in \mathbf{Z}$) colliding at crossing a rectilinear topologically nontrivial thread $A_\theta(n)$.

This effect takes place actually within a fixed topological domain inside the discrete vacuum manifold R_{YM} , possessing the topological number n .

Colliding magnetic charges with different topological numbers are suppressed in the spatial region near the axis z (of the chosen rest reference frame) by large widths of domain walls in this spatial region.

The said means the possible annihilation of all the topologically nontrivial YM vacuum BPS monopole modes and excitations over this BPS monopole vacuum (suffered the Dirac fundamental quantization [4]) during a definite time.

As a consequence of such possible annihilation, Higgs (BPS monopole) modes become free electric fields (at least in the region out of vortices cores, where superfluid potential motions occur inside the vacuum manifold R_{YM}): their electric charges e are dual (due

to the Dirac quantization [21] of electric and magnetic charges) to zero magnetic charges can only survive upon the above-described annihilation.

Such situation when Higgs modes possess arbitrary electric charges e , while magnetic charges $\mathbf{m} \neq 0$ are confined is referred to as the *Higgs phase* in modern physical literature (see e.g. [22]).

If quarks are incorporated in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac [4], disappearing topologically nontrivial YM modes via the "colliding" mechanism [9] can cause the possibility to observe free "coloured" quarks in the spatial region near the axis z .

The said can serve as a (perhaps, enough rough) representation for the asymptotical freedom of quarks in that model.

Vice versa, in the spatial region $|\mathbf{x}| \rightarrow \infty$, where walls between different topological domains inside R_{YM} become highly thin, the infrared topological and physical confinement of quarks (in the spirit [12, 23]).

Such confinement can be ensured, for instance, by surviving, upon colliding [9] at threads A_θ , topologically trivial and, perhaps, topologically nontrivial, YM modes.

In detail, the specific of the Higgs phase in the non-Abelian model (with BPS monopoles) quantized by Dirac [4] will be us discussed in Section 5.

Also we discuss important consequences for QCD (based on the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac) of assuming the "discrete" geometry.

The thing is that the Bogomol'nyi equation (1.1) is fulfilled to within the sign before the (covariant) derivative D [11].

Herewith it can be shown (repeating the arguments [9]) that this duality in specifying the sign of the vacuum "magnetic" field $\mathbf{B}$ becomes unavoidable when there exists the (already mentioned) group G_1 of global rotations, compensating changes in the vacuum (YM-Higgs) configuration (Φ^a, A_μ^a) at rotations around the axis z of the chosen (rest) reference frame.

This causes inverting the sign before the generator [9] $h \equiv h(\Phi) \equiv \Phi/a$ (with Φ being a Higgs BPS monopole mode and $a^2 \equiv m/\sqrt{\lambda}$) of the residual $U(1)$ gauge symmetry group in the quested Minkowskian Higgs model.

More exactly, to within an isomorphism, the elements [9]

$$\alpha(n) = (P \exp(-\int_0^{2\pi} A_\theta d\theta))^{-1} \quad (2.9)$$

(with P standing for parallel transports along appropriate curves [integration ways]) of the holonomies group [7, 9] $H \simeq U(1)$ may be constructed involving (topologically nontrivial) threads A_θ .

In this case there exist gauge transformations [9]

$$\tilde{h}(\Phi) = \alpha(2k)h(\Phi)\alpha(2k)^{-1} = -h(\Phi) \quad (2.10)$$

mapping the topological domain with a topological number k in the discrete group space H (and by that the appropriate topological domain inside the degeneration space R_{YM}) into the topological domain with the topological number $-k$.

Herewith the change in the sign of $\mathbf{B}$ in the Bogomol'nyi equation (1.1) occurs automatically at such mapping, accompanied also by changing the signs of magnetic charges $\mathbf{m}$.

Thus vacuum "magnetic" fields $\mathbf{B}$ and $-\mathbf{B}$ can be identified by means of gauge transformations (2.10), implicating (topologically nontrivial) threads A_θ , and this implies, regarding the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac, the manifest invariance of the appropriate vacuum "magnetic" energy, squared by $|\mathbf{B}|$, with respect to changes in the sign of the vacuum "magnetic" field $\mathbf{B}$.

The same gauge transformations (2.10) influence also the "electric" energy of the Minkowskian BPS monopole vacuum (suffered the Dirac fundamental quantization [4]), squared by the topological momentum P_N and given by the action functional (1.11).

As a result, the "electric" energy of this vacuum and the action functional (1.11) corresponding to this "electric" energy prove to be also invariant with respect to changes in signs of topological charges k .

Just this reasoning about the vacuum "magnetic" and "electric" energies of the Minkowskian BPS monopole vacuum allows to draw the conclusion that the look of the initial and residual gauge symmetries groups in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac would be modified: respectively,

$$G_M = G_0 \otimes \mathbf{Z}/\mathbb{Z}_2; \quad U_1 = U_0 \otimes \mathbf{Z}/\mathbb{Z}_2.$$

This implies the modification

$$R'_{YM} = (\mathbf{Z}/\mathbb{Z}_2) \otimes G_0/U_0$$

in the look of the appropriate degeneration space (vacuum manifold) R_{YM} .

Section 5 of the present study we devote to the specific of the Higgs phase in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac.

Above, this specific was us characterized; so we shall not repeat ourselves. On the other hand, note the following.

It seems to be a very useful thing, to compare (as it will be done in Section 5) the view about the Higgs phase at applying the Dirac fundamental quantization method [4] to the Minkowskian Higgs model (with vacuum BPS monopoles) with that associated with the so-called *dual superconducting picture* of confinement for quark and magnetic monopoles [22].

This conception of the dual superconducting picture, enveloping QCD based on the Minkowskian Higgs model with violating the "initial" $SU(2)$ gauge symmetry, includes the "Mandelstam" linearly increasing potential $O(Kr)$ (with $K = \text{const}$ being the string tension [24]) and fixing the *maximal Abelian gauge* (MAG), coming to the maximal diagonalization of the $SU(2)$ gauge matrices.

MAG can be achieved in the Yang-Mills (YM) theory by means maximizing the integral

$$\int d^4x [(A_\mu^1)^2 + (A_\mu^2)^2]; \quad A_\mu = A_\mu^a \tau_a.$$

It is equivalent to the fact (see e.g. Ref. [25]) of maximization of the quantity

$$\sum_x \sum_{\mu=1}^4 \text{Tr}[\tau_3 U_\mu(x) \tau_3 U_\mu^\dagger(x)],$$

where $U_\mu(x)$ are $SU(2)$ gauge matrices.

MAG can be got [26] by imposing the "Lorentz" gauge

$$D_\mu^{ab} A_\mu^b = 0.$$

The dual superconducting picture of confinement includes the Higgs phase, in which Higgs modes are considered [22] as "free" particles with nonzero electric charges while all the magnetic charges are confined, and also the *confinement phase* [22], dual to the Higgs phase, when Higgs modes become free and purely "magnetic" objects while gluons, monopoles and (incorporated) quarks are confined by (infinitely narrow) Meissner flux tubes [24], of the hadronic size ~ 1 fm (~ 5 GeV $^{-1}$), similar to ones in a superconductor.

Herewith the role of the Mandelstam linearly increasing potential $O(Kr)$ [24] between (electric, magnetic) charges comes in this confinement mechanism just to breaking Meissner flux tubes (strings) exceeding the hadronic size ~ 1 fm. Such breaking is advantageous from the energetic standpoint.

It is transparent, in the light of the said above, that this dual superconducting picture of confinement foresees violating the "initial" $SU(2)$ gauge symmetry group down to its $U(1)$ subgroup, that implicates the appearance of Higgs modes. In this case MAG fixing seems to be quite correct.

Indeed, it is not enough to violate the "initial" $SU(2)$ gauge symmetry group down to its $U(1)$ subgroup in order to achieve the dual superconducting picture of confinement.

On the contrary, this Abelian $U(1)$ gauge symmetry, with Higgs modes appearing upon MAG fixing, implying the breakdown of the initial $SU(2)$ gauge symmetry, would be violated farther, and this is the thing necessary for justifying the dual superconducting picture of confinement [22] in the YM theory.

Really, explaining the dual superconducting picture of confinement [22] can be found in the framework of the *Nielsen-Olesen* (NO) Abelian Higgs model [27], involving *NO vortices*.

This model serves as a highly correct representation for the Abelian Higgs theory, in which the initial $U(1)$ gauge symmetry group is violated.

On the other hand, just MAG fixing can reduce the initial non-Abelian Higgs model to the NO one.

In this case there exist specific solutions to the equations of motions, so-called NO vortices [27].

Herewith there is a highly transparent analogy between NO vortices and above-mentioned Meissner flux tubes.

Discussing briefly the NO model [27] in Section 5, we demonstrate that NO vortices are the particular case of thread topological defects inside the appropriate vacuum manifold in this model swept by the nonzero value $|\phi| \neq 0$ for the Higgs field ϕ in the "asymmetric" phase.

The important task in this situation is to ascertain the look of this manifold.

This task is connected closely with the thermodynamic specific of the Abelian Higgs model [27], involving NO vortices.

It turns out that the first-order phase transition takes place in this model, accompanied by gradual violating the $U(1)$ gauge symmetry (contrary to "momentary" violating gauge symmetries groups in theories involving second-order phase transitions [6, 28]).

The typical picture of first-order phase transitions in gauge theories is following.

At first-order phase transitions in such theories, "bubbles" [28] of "asymmetric" thermodynamic phases are "conceived" inside "symmetric" thermodynamic phases at temperatures below a Curie point T_c .

There are always prolonged processes at which "symmetric" thermodynamic phases become *metastable*, coexisting during a definite time with *stable* ("asymmetric") thermodynamic phases.

On the other hand, order parameters (a $\phi(T)$) appear (disappear) rapidly at first-order phase transitions, and this implies the finite difference ΔF of the free energy. This difference is called *the latent heat* of the given first-order phase transition [29]. Such situation coexisting two thermodynamic phases below $T = T_c$ is referred to as *supercooling* in modern literature (e.g. [29]).

When the temperature T decreases further, down to some its value $T_b < T_c$ (or when the system of fields is perturbed by either external or internal fluctuations), it drops rapidly in the "true vacuum" corresponding to the stable ("asymmetric") thermodynamic phase with $\phi(T) = \phi_0$ (herewith the system leaves the "false vacuum" $\phi(T) = 0$ corresponding to the metastable, "symmetric" thermodynamic phase).

As a result, the metastable thermodynamic phase disappears entirely, and the appropriate latent heat is liberated in this dropping process. This phenomenon is referred to as *reheating* (see e.g. [29])³.

The Abelian Higgs model [27], involving NO vortices, is the particular case of gauge theories involving first-order phase transitions.

Thus all the us discussed above remains valid also for the model [27].

The specific only in choosing the vacuum expectation value $\langle |\phi|^2 \rangle$ as the order parameter in that model, with all ensuing consequences.

³The above discussed dropping the system of fields in the "true vacuum" $\phi(T) = \phi_0$ in gauge theories implicating first-order phase transitions can be explained [29] by thermal fluctuations or by various tunnelling effects shifting this system of fields over the potential barrier (discontinuity) between $\phi(T) = 0$ and $\phi(T) = \phi_0$.

From the geometrical standpoint, the first-order phase transition in the Abelian Higgs model [27] can be explained by assuming the gradual destruction of the continuous $U(1) \simeq S^1$ group space in that model to a discrete set, can be obtained at cutting off this group space by its topological domains (there are, indeed, the countable number $\mathbf{Z} = \pi_1 S^1$ of such topological domains).

Then at the first-order phase transition taking place in the NO model [27], the "bubble" of the "asymmetric" (stable) thermodynamic phase inside the "symmetric" (metastable) one is represented just by the above discrete set.

In turn, the metastable thermodynamic phase in the NO model [27] can be represented by the continuous set consisting of a part of topological domains remained still "gluing together".

In the temperature interval $[T_b; T_c]$, corresponding to the supercooling situation [29] in the Abelian Higgs model [27], these (continuous and discrete) sets coexist.

But when the temperature descends below T_b , the metastable thermodynamic phase, represented by the topological domains "gluing together", disappears entirely. Herewith only the discrete set, corresponding to the stable thermodynamic phase and "true" vacuum $|\phi| = \phi_0$, survives.

Such is the outline of the first-order phase transition occurring in the Abelian Higgs model [27] with NO vortices.

Assuming this scheme, involving the "discrete" vacuum geometry (it can be shown that the vacuum manifold in this case is just isomorphic to the above discussed discrete set, we shall denote as $\tilde{U}$), for the first-order phase transition can serve for justifying namely NO vortices, as a particular case of thread topological defects, in the model [27].

Geometrically, NO vortices are located inside the appropriate vacuum manifold that is isomorphic to $\tilde{U}$.

The reasoning here similar to that in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac, that is the basic topic of the present study.

The distinction only in some (although greatly important) details regarding the nature of vortices in the enumerated models.

For understanding the role of the Abelian Higgs model [27] with NO vortices in the dual superconducting picture of confinement in QCD, the transparent analogy between physical processes occurring in the former model and in Type II superconductors proves to be very helpful.

Type II superconductors are described correctly in the framework of the Ginzburg-Landau model [30].

The key points in describing Type II superconductors are the following.

Firstly, the Ginzburg-Landau model [30] implicates the initial $U(1)$ gauge symmetry group, like the NO model [27]. Thus both the models are patterns of Abelian theories.

In this case the Ginzburg-Landau model [30] can be got formally from the NO model [27] (as a typical pattern of Abelian Higgs theories) by identifying a Cooper pair with the Higgs field squared $|\phi|^2$.

Secondly, it is the general property of superconductors (of Type I as well of Type II), exponential falling down [30] the magnetic field $\mathbf{B}$ at moving deep into a superconducting region:

$$\mathbf{B}(x) = \mathbf{B}_0 e^{-x/\lambda_L},$$

with

$$\lambda_L = \frac{1}{M_\gamma}$$

being the penetration or London depth, depending on the "photon mass"

$$M_\gamma = \sqrt{2} ea, \quad a \equiv |\phi|,$$

where ϕ is the wave function of the studied superconductor.

As in the NO model [27], the value a^2 plays the role of the order parameter in the Ginzburg-Landau superconductivity model [30]. $a^2 = 0$ in the symmetric (*normal*) thermodynamic phase, while $a^2 \neq 0$ in the asymmetric (superconducting) thermodynamic phase.

Thirdly, there is the following specific [13] of Type II superconductors.

Let us place a superconductor in an external magnetic field $\mathbf{H}$.

Then, in definition [13], Type II superconductors are those in which the *Ginzburg-Landau parameter* [30]

$$\kappa = \frac{\lambda_L}{\xi} < 1/\sqrt{2}.$$

In this Eq. ξ is the *coherence length*, whose physical sense is [13] the correlations radius of fluctuations in the appropriate order parameter $\langle |\phi|^2 \rangle$. Indeed, κ is a function of the temperature T : $\kappa \equiv \kappa(T)$.

It turns out [13, 30] that the inequality $\kappa < 1/\sqrt{2}$ proper to Type II superconductors is associated with the negative surface tension

$$\alpha_{ns} = \int_{-\infty}^{\infty} (f_s - f_n) dx < 0$$

between the normal (n) and superconductor (s) phases.

In the latter case it becomes advantageous, from the thermodynamic standpoint, to compensate increasing the volume energy by this negative surface energy $\alpha_{ns} < 0$.

This implies arising germs of the n-phase inside the s one at the values H of the background magnetic field H exceeding a field H_{c1} , called [13] the lower critical field. Vice versa, germs of the s-phase inside the n one arise at the values H don't exceeding a field H_{c2} , called [13] the upper critical field.

Thus in the interval $H_{c1} < H < H_{c2}$, one can observe [13] the mixed state in Type II superconductors (there are definite alloyed metals and combinations of metals). In this case a Type II superconductor abides simultaneously in the n and s states.

At $H \leq H_{c1}$, a specimen is purely in the s-state, while it is purely in the n-state at $H \geq H_{c2}$.

It is easy to see the transparent parallel between the mixed state in Type II superconductors and the coexistence of bubbles of the metastable and stable thermodynamic phases in the temperature interval $[T_b; T_c]$ in the NO model [27] (indeed, one would replace temperatures T with magnetic tensions H in the case of Type II superconductors). The mixed state in Type II superconductors is also the supercooling case [29].

Ascertaining the explicit dependence $H(T)$ (as it was performed in Ref. [13]) seems to be helpful for understanding this parallel.

Thus the first-order phase transition occurs in Type II superconductors. As a result, the conclusion can be drawn that the case of Type II superconductors can be treated as a nonrelativistic limit of the Abelian Higgs NO model [27].

As in the NO model [27], the nontrivial topological content of the Ginzburg-Landau superconductivity theory [30] can be discovered. We demonstrate that it is associated with including the electromagnetic vector potential $\mathbf{A}$ in the expression in [13]

$$\mathbf{j}_s = \frac{\hbar e}{2m} n_s (\nabla \phi - \frac{2e}{\hbar c} \mathbf{A})$$

for the gauge ($U(1)$) invariant superconductivity current $\mathbf{j}_s$ (involving the density n_s of superconductive electrons).

In this case

$$\text{rot } \mathbf{j}_s \sim \text{rot } \mathbf{A} \neq 0,$$

and this is the sign of nontrivial topologies in the Ginzburg-Landau superconductivity theory [30] according the arguments similar to those explained topologically nontrivial vortices [15] in a liquid helium II specimen (at rest).

More exactly, since

$$\oint_C \mathbf{A} d\mathbf{l} = \int \text{rot } \mathbf{A} \cdot d\mathbf{f} = \int \mathbf{B} d\mathbf{f} \equiv \Phi,$$

where in this context, Φ is the flux of the magnetic field $\mathbf{B}$,

$$\oint_C \mathbf{A} d\mathbf{l} = \frac{\hbar c}{2e} \oint_C \nabla \Phi \cdot d\mathbf{l} = \frac{\hbar c}{2e} \delta \Phi$$

for an integration way C lying inside a superconductor far from its surface.

However from the claim for the superconductor wave function to be finite, it follows that this change of the phase Φ can take only values $2\pi n$ ($n \in \mathbf{Z}$).

One comes thus to the result [13]

$$\Phi = n\Phi_0, \quad \Phi_0 = \frac{\pi \hbar c}{|e|} = 2 \cdot 10^{-7} \text{ G cm}^2$$

for the phase Φ . The value Φ_0 is called *the elementary quant of the magnetic flux* [13].

The variation $\delta\Phi$ is indeed infinitesimal, that corresponds to the fact that $\text{rot } \nabla\Phi = 0$. Thus one can observe actual disappearing nontrivial topologies deep into a superconductor.

Indeed, in Type II superconductors the background magnetic field H penetrates into the superconducting specimen along the parallel each other lines, so-called *Abrikosov threads*, forming a lattice. These serve as embryos of the normal phase in the specimen, generating simultaneously circular superconducting currents inside it [13]. Note that these Abrikosov threads belonging to the “continuous” thermodynamical phase n are additional to those would exist inside the discrete vacuum manifold $\tilde{U}(1)$.

In the present study Abrikosov threads will be analyzed repeating the arguments [13]. In particular, it will be argued that the first order phase transition occurring in Type II superconductors can be reduced to the non-smooth behaviour of the order parameter $|\phi(r)|^2$ in the points locating Abrikosov threads.

The geometric (topological) specific of that first order phase transition will be also discussed and the series of hypotheses concerning this specific will be suggested.

In the light of the said about the Abelian Higgs NO model [27] involving NO vortices and a parallel between this and the Ginzburg-Landau theory [30], especially in that part of the latter one concerning Type II superconductors, the dual superconducting picture of confinement in QCD can be explained enough good as that corresponding to the stable thermodynamic phase, below the temperature T_b (when the metastable phase disappears entirely and the system fall down in the “true” vacuum $|\phi| = \phi_0$), in the NO model [27].

In this stable thermodynamic phase, an external magnetic field $\mathbf{B}$, being present, is concentrated [31] in cores of vortices and will not extend into the superconducting region.

As a result, the interaction energy of magnetic monopoles grows linearly with their separation d [31]:

$$V = \frac{1}{2} \int d^3x \mathbf{B}^2 \propto \frac{4\pi d}{e^2 \lambda_L^2},$$

as it would be in the dual superconducting picture [22] of confinement.

In the modern literature the above discussed (phenomenological) picture of confinement based on MAG fixing and the dual superconducting picture (similar to that one discovers in the Ginzburg-Landau model [30] concerning Type II superconductors and implicating [22] the Higgs and confinement phases) is referred to as the *Abelian dominance* [32]).

The situation changes rather in a radical way, in comparison with the Abelian dominance [32] picture of confinement, in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac [4].

There are two important “points of distinction” between the “Abelian dominance” [32] sight on confinement and that sight based upon the Dirac fundamental quantization method [4] applied to (Minkowskian) QCD and coming to the Gauss-shell reduction of the appropriate Hamiltonian.

Firstly, at constructing Minkowskian Gauss-shell QCD (underlying by vacuum BPS monopole solutions) the specific gauge, different from MAG, can be fixed in the more general case when the initial $SU(3)_{\text{col}}$ gauge symmetry group inherent in Minkowskian Gauss-shell QCD is violated in the

$$SU(3)_{\text{col}} \rightarrow SU(2)_{\text{col}} \rightarrow U(1) \quad (2.11)$$

wise.

Then interesting physical effects can be achieved at choosing the antisymmetric Gell-Mann matrices $\lambda_2, \lambda_5, \lambda_7$ as generators for $SU(2)_{\text{col}}$ [1, 5, 12].

Among such consequences fixing the ("antisymmetric") "gauge", one can mention fermionic rotary (axial) degrees of freedom $\mathbf{v}_1 = \mathbf{r} \times \mathbf{K}$ (with $\mathbf{K}$ being the polar colour vector, $SU(2)_{\text{col}}$ triplet).

These fermionic rotary degrees of freedom $\mathbf{v}_1$ somewhat similar to rotary terms one can detect in molecules.

On the other hand, to explain the appearance of above fermionic rotary degrees, assuming the "discrete" group geometry for $SU(2)_{\text{col}}$ and $U(1)$ in Eq. (2.11) (then also the appropriate vacuum manifold $SU(2)_{\text{col}}/U(1)$ possesses a "discrete" geometry) seems to be plausible.

Secondly, the Gauss-shell reduction of Minkowskian QCD in the BPS monopole background implies [1, 5, 7, 8, 12, 20] the Green function of the Gauss law constraint can be expand by the Coulomb and "golden section" potentials.

Thus the role of the Mandelstam linearly increasing potential $\sim Kr$ (and the dual superconducting picture [22, 31] together with) in Minkowskian constraint-shell QCD is underestimated considerably, at least at low temperatures $T \rightarrow 0$.

And only at enough high temperatures $T \neq 0$, a chance appears to incorporate the Mandelstam linearly increasing potential in Minkowskian constraint-shell QCD with vacuum BPS monopoles.

Such possibility will be also us discussed in Section 5.

3 How one can introduce "discrete" geometry for de-generation spaces in Minkowskian Higgs models.

Recall again the recent paper [6]. In this paper some modern Minkowskian Higgs models with *stationary* vacuum monopole solutions were analysed.

Besides already mentioned BPS monopole model [7, 8, 9, 10, 11], one can mention the 't Hooft-Polyakov monopole model [33, 34].

As it was discussed in [6], such (vacuum) monopole solutions are compatible with the "continuous" vacuum geometry

$$R \equiv SU(2)/U(1) \simeq S^2. \quad (3.1)$$

Also it was argued that Minkowskian Higgs models with stationary vacuum monopole solutions, involving the "continuous" vacuum geometry (3.1), can be described quite correctly in the framework of the Faddeev-Popov (FP) "heuristic" quantization scheme [35], coming [1, 6] to actual fixing the Weyl gauge $A_0 = 0$ (for instance via the multiplier $\delta(A_0)$ in appropriate FP path integrals).

From the topological viewpoint, the *continuous* vacuum manifold R ⁴, (3.1), possesses the one kind of topological defects, namely the point hedgehog topological defects.

These topological defects are associated with vacuum monopole solutions in appropriate Minkowskian Higgs models.

Point hedgehog topological defects inside R are determined by the topological chain [6, 7]

$$\pi_2(R) \equiv \pi_2 S^2 = \pi_1(H) \equiv \pi_1 U(1) = \mathbf{Z}. \quad (3.2)$$

From the thermodynamic points of view, all the kinds of topological defects may be explained (repeating the arguments [9]) by violating the thermodynamic equilibrium over a region in the appropriate coordinate (in particular, Minkowski) space.

For example, point hedgehog topological defects inside R are associated with violating the thermodynamic equilibrium in an infinitesimal neighbourhood of the origin of coordinates. Such neighbourhoods are topologically equivalent to the two-sphere S^2 .

In the present study we attempt to demonstrate our readers that going out from the FP "heuristic" quantization scheme [35] to the Dirac "fundamental" one [4] in the Minkowskian Higgs model with vacuum BPS monopole solutions claims the in principle new geometrical approach to constructing the appropriate vacuum manifold (in comparison with assuming the "continuous" $\sim S^2$ vacuum geometry in the former case).

This "new geometrical approach" (whose outlines were contemplated in the recent paper [1]) comes to assuming the "discrete" geometry for the vacuum manifold $SU(2)/U(1)$.

As we shall make sure soon, this assuming explains enough good various vacuum rotary effects [3] inherent in the physical BPS monopole vacuum suffered the Dirac fundamental quantization [4].

⁴Following [9], R may be called the *degeneration space*.

It is quite correct herewith to interpret the initial symmetry group G in a Minkowskian Higgs model as that does not change the energy functional (Hamiltonian) of that model, while the residual symmetry group $H \subset G$ as that consisting of transformations that keep invariant a fixed equilibrium state.

All these states (at a fixed temperature $T < T_c$, with T_c being the appropriate Curie point in which the initial symmetry G is violated and the second-order phase transition occurs) just form the appropriate degeneration space $R = G/H$. The natural claim to this space is herewith *to be topological*.

The structure of a degeneration space may be investigated with the aid of the Landau theory of second-order phase transitions.

An equilibrium state is determined by the condition for the free energy of the given system to be minimal.

In the Landau theory of second-order phase transitions (the pattern of which is the Minkowskian Higgs model) one supposes that an equilibrium state may be found at minimizing the free energy of the given system by the set of states specified by a finite number of parameters (called *order parameters*), but not by the set of all the states.

So, let us represent [1] the initial, $SU(2) \equiv G$, and residual, $U(1) \equiv H$, gauge symmetries groups in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac in the shape of discrete spaces

$$SU(2) \simeq G_0 \otimes \mathbf{Z}; \quad U(1) \simeq U_0 \otimes \mathbf{Z}, \quad (3.3)$$

respectively ⁵.

From the topological viewpoint, the discrete representation (3.3) for the gauge groups G and H extracts "small" (topologically trivial) and "large" (corresponding to topological numbers $n \neq 0$) gauge transformations in the complete set of appropriate gauge transformations (the idea of such subdividing for gauge transformations was suggested in Ref. [18]).

According to the terminology [18], the complete groups G_0 and H_0 just contain "small" gauge transformations, that implies

$$\pi_n G_0 = \pi_n H_0 = 0 \quad (3.4)$$

for loops in the group spaces G_0 and H_0 in all the dimensions $n \geq 1$.

Simultaneously, in definition,

$$\pi_0 G_0 = \pi_0 H_0 = 0, \quad (3.5)$$

i.e. G_0 and H_0 are *maximal connected components* [9] in their gauge groups (respectively, G and H).

Later Eq. implies [9] that

$$\pi_0[G_0 \otimes \mathbf{Z}] = \pi_0[G_0 \otimes \mathbf{Z}] = \pi_0(\mathbf{Z}) = \mathbf{Z}. \quad (3.6)$$

The latter relation indicates transparently the discrete nature of the appropriate group spaces ⁶.

To construct the degeneration space (vacuum manifold) $R_{YM} \equiv G/H$ corresponding to the discrete representation (3.3) for the gauge groups G and H , let us multiply Eq. [6, 36]

$$G = H \oplus G/H \quad (3.7)$$

left by $\mathbf{Z}$, substituting simultaneously in (3.7) the "discrete" representations (3.3) for the gauge symmetries groups:

$$\mathbf{Z} \otimes G_0 = \mathbf{Z} \otimes U_0 + \mathbf{Z} \otimes G_0/U_0. \quad (3.8)$$

⁵Such representation for gauge groups spaces was proposed for the first time in the paper [16].

⁶The origin of Eq. (3.6) was revealed in the monograph [9], in §T20.

It is the particular case of the more general relation

$$\pi_i(K) = \pi_i(L_1) + \dots + \pi_i(L_r)$$

for a group K which is the product of the groups $L_1 \dots L_r$ at a fixed i (it is correctly for the Lie groups of the series SU , U and SO , with which modern theoretical physics deals).

From latter Eq. we learn that the degeneration space in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac [4] is

$$R_{\text{YM}} = \mathbf{Z} \otimes G_0/U_0. \quad (3.9)$$

Again $\pi_1(R_{\text{YM}}) = \mathbf{Z}$ since $U_0 \otimes \mathbf{Z}$ is multi-connected.

It is correctly because the isomorphism (2.1) (the proof of this isomorphism was given in Ref. [9], in §T20, we recommend our readers for understanding the matter).

Further, it becomes transparent from Eq. (3.9) that the "small" coset G_0/U_0 is one-connected:

$$\pi_1(G_0/U_0) = 0.$$

Really, the coset G_0/U_0 is treated as the space of U_0 -orbits on G_0 ; the latter space is one-connected.

One can see also the topological equivalence between G_0/U_0 and the subset of one-dimensional ways on R_{YM} can be contracted into a point.

The vacuum manifold R_{YM} is transparently multi-connected:

$$\pi_0(R_{\text{YM}}) = \mathbf{Z}. \quad (3.10)$$

This Eq. may be explained in the framework of our above remark [9] about homotopical groups of products of groups.

Latter Eq. has very important physical sense.

It implies [9] that domain walls exist between different topological sectors in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac.

The origin of said domain walls is in the "discrete" factorisation (3.3) of the residual gauge symmetry group $U(1)$ ⁷.

⁷Indeed, the above-mentioned "discrete" factorisation (3.3) of the residual gauge symmetry group $U(1)$ can be referred to as a "discrete-continuous" in contrast to purely "discrete" symmetries.

The typical look of latter symmetries is [37] $\exp(2\pi i/p)$ ($p \in \mathbf{Z}$). If, for instance, a single complex scalar field ς is given, the appropriate low energy theory will be invariant under the transformation

$$\varsigma \rightarrow \exp(2\pi i/p) \varsigma.$$

On the other hand, as it was noted by Krauss and Wilczek [37], the most important consequence of continuous local (gauge) symmetries is the existence of gauge fields. Such fields are introduced in order to formulate covariant derivatives. These are necessary since invariant interactions involving gradients can be formed; these interactions, in turn, are necessary in order that charged fields can propagate. In the case of a discrete symmetry there is no similar need to introduce a gauge potential because the ordinary derivative already transforms simply.

Thus one can assert that the distinction between the original gauge low energy theory and the "discrete" model is in the absence of the interaction items in the latter case.

The on principle another situation is in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac, the subject of the present discussion.

In this model the factorisation (3.3) (calling for justifying the Dirac quantization scheme [4] applying here), the presence of the "continuous" factors U_0 and G_0 in Eqs. (3.3) and (3.9) comes off the Krauss-Wilczek prohibition [37] for interactions.

As it is well known (see e.g. §7.2 in [28] or the paper [19]), the width of a domain (or *Bloch*, in the terminology [19]) wall is roughly proportional to the inverse of the lowest mass of all the physical particles in the (gauge) model considered.

In Minkowskian Higgs models the typical such scale is the (effective) Higgs mass $m/\sqrt{\lambda}$.

In particular, in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac [4], $m/\sqrt{\lambda}$ is the only mass scale different from zero.

This fact was explained in Ref. [2]. This was argued by zero masses of YM fields represented by topological Dirac variables A_i^D ($i = 1, 2$).

These variables become massless at the Dirac removal [2, 5]

$$U(t, \mathbf{x})(A_0^{(0)} + \partial_0)U^{-1}(t, \mathbf{x}) = 0. \quad (3.11)$$

of temporal YM components A_0 , that are, indeed, nondynamical fields possessing zero canonical momenta

$$E_0 \equiv \partial \mathcal{L} / \partial (\partial_0 A_0) = 0.$$

In this context [7, 8],

$$U(t, \mathbf{x}) = v(\mathbf{x}) T \exp \left\{ \int_{t_0}^t \left[\frac{1}{D^2(\Phi^{\text{BPS}})} \partial_0 D_k(\Phi^{\text{BPS}}) \hat{A}^k \right] d\bar{t} \right\}, \quad (3.12)$$

It is so since now in each topological domain of the vacuum manifold R_{YM} , (3.9), two “large” (stationary) gauge matrices corresponding to a fixed topological number n can be always connected by a “small” gauge transformation. This can be written down as [5]

$$v^{(n)}(\mathbf{x}) \rightarrow v^{(n)'}(\mathbf{x}) = v^{(n)}(\mathbf{x}) u_0^{-1}; \quad u_0 \in U_0.$$

Appropriately, gauge fields setting “large” gauge matrices $v^{(n)}(\mathbf{x})$ are transformed according to the law

$$\frac{\sigma \cdot \mathbf{A}_\mu^{(n)}}{2} \rightarrow \frac{\sigma \cdot \mathbf{A}_\mu^{(n)'}}{2} = u_0 \left(\frac{\sigma \cdot \mathbf{A}_\mu^{(n)}}{2} \right) u_0^{-1} - \frac{i}{g} [\partial_\mu u_0^{-1}] u_0;$$

with σ being Pauli matrices and g being the (YM) coupling constant.

This transformation law remains correct, in particular, for topological Dirac variables A_i^D ($i = 1, 2$) [5, 20] (see below), which, by their construction [12] are gauge invariants.

Thus in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac, with the “discrete-continuous” vacuum geometry (3.9) been assumed, interactions “have vested right to existence”.

This permission for interactions has a crucial importance for the mentioned model. For example, in the BPS (Wu-Yang) monopole background there exist the such solution for the Gauss law constraint permitting the expansion [5, 7, 8, 12, 20] in terms of the Coulomb and “golden section” potentials. This, in turn, serves as base for describing the four-fermionic instantaneous interaction. In the quark sector of the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac this is the initial step in constructing quarkonium ($q\Gamma\bar{q}$ with Γ being a combination of Dirac matrices) bound states. The algorithm constructing such bound states utilizing the Markov-Yukawa prescription [38] (see e.g. §3.1 in the revue [12]).

And moreover, the mentioned “golden section” instantaneous potential can lead [12] to rearrangement of the naive perturbation series of the spontaneous chiral symmetry breaking type (especially, in the infrared momentum region).

with $v(\mathbf{x})$ being stationary Gribov topological multipliers (depending indeed on topological numbers n).

If, additionally, the transverse gauge (1.7) [5] is fixed for topological Dirac variables A_i^D , this just implies the masslessness of YM fields. Here is a transparent analogy with the photon case in QED. In that case one gets plane waves by fixing the Lorentz gauge

$$\phi = 0; \quad \partial_i A^i = 0 \quad (i = 1, 2, 3) \Leftrightarrow \partial_\mu A^\mu = 0$$

(with $A_0 \equiv \phi$ being the scalar potential).

As it was argued in the monograph [39], such fixing the (relativistic invariant) Lorentz gauge implies the equation of motion (referred to as the *D'alambert* or *wave* equation)

$$\Delta \mathbf{A} - \frac{\partial^2 \mathbf{A}}{\partial t^2} = 0 \quad (3.13)$$

(at setting $c = 1$ for the light velocity). Latter Eq. shows transparently the absence of the (rest) mass for a photon.

As it was stressed in Ref. [3] and in Section 2 of the present work, in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac [4], the typical value of the length dimension inversely proportional to $m/\sqrt{\lambda}$ is the size ϵ of BPS monopoles.

It may be given as [5, 7, 8]

$$\frac{1}{\epsilon} = \frac{gm}{\sqrt{\lambda}} \sim \frac{g^2 \langle B^2 \rangle V}{4\pi}. \quad (3.14)$$

Thus ϵ is, in turn, inversely proportional to the spatial volume $V \sim r^3$ occupied by the appropriate (YM-Higgs) field configuration.

The said allows to assert that ϵ disappears in the infinite spatial volume limit $V \rightarrow \infty$, while it is maximal at the origin of coordinates (herewith it can be set $\epsilon(0) \rightarrow \infty$).

This means, due to the above reasoning [19], that walls between topological domains inside R_{YM} become infinitely wide, $O(\epsilon(0)) \rightarrow \infty$, at the origin of coordinates.

The fact $\epsilon(\infty) \rightarrow 0$ is also meaningful. This implies actual merging of topological domains inside the vacuum manifold R_{YM} , (3.9), at the spatial infinity.

This promotes the infrared topological confinement (destructive interference) of Gribov "large" multipliers $v^{(n)}(\mathbf{x})$ in gluonic and quark Green functions in all the orders of the perturbation theory.

The latter fact was demonstrated utilizing the strict mathematical language in Ref. [23] (partially these arguments [23] were reproduced in [1]).

The existence of domain walls between different topological sectors of the vacuum manifold R_{YM} , (3.9), permits their treatment as *quite separated sets*.

The notion of quite separated sets [40] means in this case that always there exists such a function $f : R_{YM} \rightarrow I$ ($I \equiv [0, 1]$) for two homotopical classes A and B inside R_{YM} that $f(x) = 0$ as $x \in A$, while $f(x) = 1$ as $x \in B$. Here x is a (one-dimensional) way.

One speaks in this case that f *separates* the sets A and B .

The nontrivial isomorphism (2.1) [9] satisfying (as we have discussed above) for the vacuum manifold R_{YM} , (3.9), implies the presence of thread topological defects inside this manifold.

On the level of phenomenology this implies [9] violating the thermodynamic equilibrium along definite lines in the given vacuum manifold.

In the simple case when such lines, threads, are straight, the thermodynamic equilibrium is violated in an infinitely narrow tube around each of these threads. Thus regions of thread topological defects have the topology of a cylinder (topologically equivalent to the circle S^1). But it is well known that $\pi_1 S^1 = \mathbf{Z}$.

In our researches about the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac [4] we assume that the region of thread topological defects inside the vacuum manifold R_{YM} , (3.9), is the infinitely narrow tube of the infinite length around the axis z in the chosen rest reference frame. The effective "diameter" of this cylinder of the infinite length is just $O[\epsilon(\infty)]$, that is, indeed, an infinitely small value.

In the next section we shall ground the above assumption investigating the properties [9] of rectilinear threads A_θ inside R_{YM} .

Furthermore, it may be shown that

$$\pi_2(R_{YM}) = \mathbf{Z} = \pi_1 H. \quad (3.15)$$

The isomorphism $\pi_1(H) = \mathbf{Z}$ follows from the arguments of the holonomies group, stated in Refs. [7, 9].

Actually, these come to the existence of one-dimensional loops in the H group space, forming holonomies elements

$$b_\gamma = P \exp(-\oint_\Sigma \mathbf{T} \cdot A_\mu dx^\mu) \quad (3.16)$$

over one-dimensional cycles Σ .

The holonomies elements b_γ , belonging to the $U(1) \subset SU(2)$ embedding, make the complete holonomies group (we shall denote all as H) isomorphic to $U(1)$ [7].

Eq. (3.16) makes the isomorphism $\pi_1(H) = \mathbf{Z}$ highly transparent (as that coming to $\pi_1 S^1 = \mathbf{Z}$).

The relation

$$\pi_2(R_{YM}) = \pi_1 H,$$

is the particular case of more general Eq. (2.2).

The good proof of the isomorphism (2.2) was demonstrated in the monograph [9], in §T20, and we recommend our readers this monograph to acquaint with it.

The nontrivial isomorphism (3.15) testifies in favour of point hedgehog topological defects inside the vacuum manifold R_{YM} in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac [4].

These point hedgehog topological defects and vacuum BPS monopole solutions are indivisible.

As in the case of (vacuum) 't Hooft-Polyakov monopole solutions [33, 34], associated with the "continuous" vacuum $\sim S^2$ geometry (3.1) and the FP "heuristic" quantization scheme [35], point hedgehog topological defects inside the discrete vacuum manifold R_{YM} , (3.9), come to violating the thermodynamic equilibrium in an infinitesimal neighbourhood of the origin of coordinates.

In particular, this violating implies the singularity of the appropriate vacuum "magnetic" fields $\mathbf{B}$, serving as the order parameters in the mentioned Higgs models, at the origin of coordinates.

In spite of definite distinctions between the 't Hooft-Polyakov and BPS Higgs models (these distinctions were pointed out, for instance, in the recent paper [6]; the principal of them is the absence of superfluid properties in the 't Hooft-Polyakov model [33, 34]) the vacuum "magnetic" field $\mathbf{B}$ diverges as r^{-2} at the origin of coordinates in both models ⁸:

$$B_k \sim \frac{r_k}{gr^3}. \quad (3.17)$$

Moreover, there is a likeness between Eqs. (3.15) and (3.2) as the particular cases of the general relation (2.2) [9], inducing point topological defects in (gauge) theories.

On the other hand, there are no thread topological defects inside the continuous vacuum manifold R , (3.1), since $\pi_1 S^2 = 0$ and moreover when the "continuous" geometry is assumed for the residual $U(1)$ gauge symmetry in the (Minkowskian) Higgs model, also $\pi_0 U(1) = 0$.

In the next section it will be argued that the first-order phase transition occurs indeed in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac [4].

To facilitate this job, now we should like discuss an useful analogy, us give the liquid helium II model.

Just in the latter model, possessing manifest superfluid properties [14], also the first-order phase transition happens.

It is associated with (spontaneous) violating the superfluidity [14] in a (resting) liquid helium II specimen along definite (rectilinear) lines, threads.

This implies [15] the presence of thread (rectilinear) vortices in that specimen.

The similar situation with arising (rectilinear) vortices takes place when this specimen turns together with the cylindrical vessel in which it is contained.

The phenomenology of the latter case was stated enough good in the monograph [13], in §29. We don't intend, in the present study, to retail all the said in the monograph [13] about this subject; however we, for all that, would like to elucidate some theses of the helium turning model. This will be very helpful for us in the next section.

A good argument in favour of the first-order phase transition occurring in the liquid helium II turning model is the nontrivial contribution ΔE [13] in the total helium (He^4)

⁸For the 't Hooft-Polyakov model [33, 34] this was demonstrated in Ref. [36], while for the BPS Higgs model, in the original papers [10].

Hamiltonian ⁹ from the global solid *potential* rotations of the helium specimen turning together with the cylindrical vessel where it is contained.

This contribution is given with Eq.

$$\Delta E \sim L\pi\rho_s \frac{\hbar^2}{m^2} \ln \frac{R}{a} : \quad (3.18)$$

with L being the length of the vessel, ρ_s being the density of the superfluid component in the helium II specimen, m being the mass of a helium atom; at last R and a are, respectively, the radius of the vessel and an arbitrary distance of atomic scales.

The shortcoming of this Eq. is its logarithmic divergence at $R/a \rightarrow \infty$.

The important peculiarity of the liquid helium II turning model [13] is the appearance of rectilinear threads (accompanied by appropriate vortices), parallel to the rotation axis

⁹ This has the typical look [41]

$$\begin{aligned} \hat{\mathcal{H}} &= \int d^3r \hat{a}^+(\mathbf{r}, t) \left(\frac{-\hbar^2}{2m} \nabla^2 \right) \hat{a}(\mathbf{r}, t) \\ &+ \frac{1}{2} \int d^3r \int d^3r' \hat{a}^+(\mathbf{r}, t) \hat{a}^+(\mathbf{r}', t) V(|\mathbf{r} - \mathbf{r}'|) \hat{a}(\mathbf{r}', t) \hat{a}(\mathbf{r}, t). \end{aligned}$$

Boson creation and annihilation operators: $a^+(\mathbf{r}, t)$ and $a^-(\mathbf{r}, t)$, respectively, can be expressed in terms of the phase $\Phi(t, \mathbf{r})$ of the Bose condensate wave function (1.4) [13].

Note that such expressing boson creation and annihilation operators in terms of the one phase function $\Phi(t, \mathbf{r})$ is in a good agreement with the quantum-mechanical calculations about the liquid helium II proposed by Bogolubov and co-authors [42], in particular, with the Bogolubov transformations [42, 43, 44]

$$\hat{b}_{\mathbf{p}}^+ = u_{\mathbf{p}} \hat{\xi}_{\mathbf{p}}^+ + v_{\mathbf{p}} \hat{\xi}_{-\mathbf{p}},$$

with

$$u_{\mathbf{p}}^2 - v_{\mathbf{p}}^2 = 1$$

and the creation (respectively, annihilation) operators

$$\hat{b}_{\mathbf{p}}^+ = \frac{\hat{a}_0 \hat{a}_{\mathbf{p}}^+}{\sqrt{n_0}}, \quad \hat{b}_{\mathbf{p}} = \frac{\hat{a}_0^+ \hat{a}_{\mathbf{p}}}{\sqrt{n_0}},$$

expressed through the "initial" creation (annihilation) operators entering actually the Bogolubov model Hamilton operator, can be represented as [42, 43]

$$\hat{H} = - \sum_{a=1}^N \frac{\hbar^2}{2m} \Delta_a + \frac{1}{2} V(|\mathbf{r}_a - \mathbf{r}_b|),$$

(with $V(|\mathbf{r}_a - \mathbf{r}_b|)$ being the interaction energy between particles a and b and N being the complete number of particles in the considered system), brought in the diagonal form

$$\hat{H} = \hat{H}_0 + \sum_{\mathbf{p} \neq 0} \epsilon(\mathbf{p}) \hat{\xi}_{\mathbf{p}}^+ \hat{\xi}_{\mathbf{p}}$$

via the above Bogolubov transformations.

Herewith the creation (respectively, annihilation) operators $\hat{a}_0^+$ and $\hat{a}_0$ corresponds to the zero momenta $\mathbf{p} = 0$ and the number n_0 of helium atoms possessing these zero momenta.

(altitude) of the cylindrical vessel (herewith chosen to coincide with the axis z of the given rest reference frame).

The liquid helium II turning model [13] possesses initially the manifest $U(1) \otimes O(2)$ symmetry.

In this model $O(2)$ is the group of *global* (rigid) potential rotations around the axis z of the liquid helium II specimen and the cylindrical vessel where it is contained.

Unlike $O(2)$, $U(1)$ is the gauge symmetry group.

In the "general" liquid helium II theory [14, 42], the latter group can be represented by matrices

$$v_n(\mathbf{r}) = \exp(i n \lambda(\mathbf{r})) \quad (3.19)$$

maintaining the Bogolubov (diagonalized) Hamiltonian $\hat{\mathcal{H}}$ [41, 42, 43].

Herewith the phase $\Phi(t, \mathbf{r})$ of the complex-value helium Bose condensate wave function $\Xi(t, \mathbf{r})$ [13], (1.4), is suffered the gauge transformations

$$\Phi(t, \mathbf{r}) \rightarrow \Phi(t, \mathbf{r}) + n \lambda(\mathbf{r}), \quad n \in \mathbf{Z}. \quad (3.20)$$

And again, "small" and "large" gauge transformations [18] can be picked out among (3.19), (3.20).

As it is well known, superfluidity phenomena [14] in a liquid helium II specimen are associated with violating the initial $U(1)$ gauge symmetry of the Bogolubov Hamiltonian $\hat{\mathcal{H}}$. This second-order phase transition occurs in the fixed Curie point $T_c \rightarrow 0$.

To explain in this case rotary effects (coming to appearance of thread vortices) in the liquid helium II turning model [13] (as well as in the liquid helium II at rest theory [15]), it is worth to assume that violating the initial $U(1)$ gauge symmetry in the liquid helium II theory in such a wise that the $U(1) \simeq S^1$ group space turns into the discrete quite separated set [40].

More exactly, it may be assumed the spontaneous breakdown

$$U(1) \simeq U_0 \otimes \mathbf{Z} \longrightarrow \mathbf{Z} \simeq \tilde{U}(1); \quad \pi_0 U_0 = \pi_1 U_0 = 0. \quad (3.21)$$

Geometrically, the discrete (quite separated) set $\tilde{U}(1)$ may be seen as the circle S^1 cut off by its topological domains (vice verse, to get the continuous $U(1)$ group space, these topological domains would be again pasted together).

On the other hand, in the liquid helium II turning model [13], the appropriate group $O(2)$ of rigid rotations remains exact.

This implies the (spontaneous) breakdown

$$O(2) \otimes U(1) \simeq O(2) \otimes U_0 \otimes \mathbf{Z} \longrightarrow O(2) \otimes \mathbf{Z} \simeq O(2) \otimes \tilde{U}(1) \equiv H \quad (3.22)$$

of the initial symmetry group $O(2) \otimes U(1)$ inherent in the liquid helium II turning model [13].

Upon some mathematical manipulations, resembling somehow ones [36] led to Eq. (3.9) in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac,

the degeneration space in the liquid helium II turning model [13] may be founded. It proves to be

$$[\mathbf{Z} \otimes O(2)] \otimes U_0 \equiv R, \quad (3.23)$$

and it is topologically equivalent to S^1 .

The latter fact may be explaining by reasoning that the group $O(2)$ of 2×2 orthogonal matrices with determinants ± 1 always may be represented as

$$SO(2) \otimes \mathbb{Z}_2,$$

with $SO(2) \simeq U(1)$ being the group of orthogonal matrices with determinants $+1$, while $\mathbb{Z}_2 \subset \mathbf{Z}$.

It is easy to see that $\pi_1 R = \mathbf{Z}$.

In detail, this may be argued by the above ascertained topological equivalence between the liquid helium II turning degeneration space R and the circle S^1 .

On the other hand, also $\pi_0 H = \mathbf{Z}$ due to the manifest presence of the discrete multiplier $\tilde{U}(1)$ in Eq. (3.22).

Thus the general isomorphism (2.1) [9] is satisfied in the case of the liquid helium II turning model [13].

In this concrete case it implies the presence of (rectilinear) thread topological defects (vortices) in the quested model.

These rectilinear vortices contribute with the item ΔE [13], (3.18), additional to the Bogolubov helium (diagonalized) Hamiltonian $\hat{\mathcal{H}}$ [41].

Just this increasing ΔE the helium energy (in comparison with that given by $\hat{\mathcal{H}}$ and referring to the helium specimen at rest) is the sign of the first order phase transition occurring.

Of course, there are walls between different topological domains inside the discrete liquid helium II turning degeneration space R , (3.23).

On the other hand, there are no point topological defects inside R since $\pi_2 S^1 = 0$.

The case when a liquid helium II is at rest is somewhat simpler than the case [13] of the liquid helium II turning together with the cylindrical vessel where it is contained.

In the former case, to justify the spontaneous appearance of rectilinear vortices ¹⁰, it may be presumed that the initial (continuous) $U(1)$ gauge symmetry group of the

¹⁰As it was explained in [15] (see also [3]), the appearance of rectilinear vortices in a liquid helium II specimen is set by Eq.

$$n = \frac{m}{2\pi\hbar} \oint_{\Gamma} \mathbf{v}^{(n)} d\mathbf{l}; \quad n \in \mathbf{Z}.$$

This Eq. implicates the helium mass m and the tangential velocity $\mathbf{v}^{(n)}$ of a rectilinear vortex.

In this case, trivial topologies $n = 0$ correspond to disappearing the cyclic integral on the right-hand side of latter Eq. It is just the case of superfluid potential motions in the given liquid helium II specimen with $\text{rot } \mathbf{v}^{(0)} = 0$.

In Ref. [3], the explanation of vortices [15] in a (rested) liquid helium II specimen as a particular case of the *Josephson effect* [17] (coming to circular persistent motions of material points) was given.

Bogolubov helium Hamiltonian $\widehat{\mathcal{H}}$ [41] is violated in the

$$U(1) \rightarrow \tilde{U}(1) \quad (3.24)$$

wise.

As in the liquid helium II turning case [13], $\pi_0 \tilde{U}(1) = \mathbf{Z}$.

Herewith the set $\tilde{U}(1)$ (as well as the set $\tilde{U}(1) \otimes O(1)$ in the case [13] of a liquid helium II specimen turning together with the cylindrical vessel where it is contained) can be treated as the residual symmetry group in the discussed helium models.

On the other hand, it can be demonstrated (applying the manipulations similar to those necessary in the liquid helium II turning case [13] to ascertain the look (3.23) for the appropriate degeneration space R) that the degeneration space $\tilde{R}$ in the liquid helium II at rest theory coincides (to within an isomorphism)¹¹ with the residual gauge symmetry group

$$\tilde{U}(1) \simeq U_0 \otimes \mathbf{Z}.$$

Thus, formally,

$$\pi_1 \tilde{R} = \pi_0 \tilde{U}(1) = \mathbf{Z},$$

and this just implies the existence of thread topological defects (rectilinear vortices) in the liquid helium II at rest theory.

The alone fact [3, 15] referring vortices in a liquid helium II at rest to nontrivial topologies $n \neq 0$, while superfluid potential motions are referred to trivial topologies $n = 0$, is very remarkable.

This allows to associate various superfluidity effects in a liquid helium II at rest specimen with "small" $U(1)$ gauge matrices while "large" ones with (rectilinear) vortices therein.

All these gauge matrices are given (to within the natural isomorphism $U(1) \simeq \tilde{U}(1)$) with Eq. (3.19).

Thus the connection between the superfluid and rotary effects in the liquid helium II model and the topological degeneration of appropriate data (the phase $\Phi(t, \mathbf{x})$ of the helium Bose condensate [13], in the first place) may be observed.

Finally, it may be argued, as in the liquid helium II turning model [13], that the first-order phase transition occurs also in the liquid helium II at rest model [15].

Returning again to the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac [4], note the very transparent analogy between the rotary item ΔE , (3.18), in the liquid helium II turning model [13] and the free rotator action functional W_N , (1.11), in the former case.

¹¹Indeed, $\tilde{R} \simeq S^1/\mathbf{Z}$. In this case, as it is easy to see,

$$\pi_1 \tilde{R} = \pi_1 S^1 = \mathbf{Z}.$$

The important distinction between the both theories is, however, in the logarithmic divergence of ΔE , (3.18), as $R/a \rightarrow \infty$, while the free rotator action functional W_N , (1.11), is suppressed actually by the value of $\epsilon(\infty) \rightarrow 0$.

The just performed brief analysis of the liquid helium II turning model [13] (as well as of the liquid helium II at rest model) suggests, by analogy, the idea to link the various rotary properties [3, 5, 7, 8, 12] of the Minkowskian Higgs model to vacuum BPS monopoles quantized by Dirac (described correctly by the free rotator action functional (1.11)) with the discrete vacuum geometry (3.9), including thread topological defects inside the appropriate vacuum manifold R_{YM} (in the similar way, has been outlined above, in which such topological defects induce rotary effects in the liquid helium II theory).

This will be the topic of the next section, where also some arguments in favour of the first-order phase transition occurring in the Minkowskian Higgs model with vacuum BPS monopole solutions quantized by Dirac and similar to that taking place in the liquid helium II theory (these arguments are connected with the "electric" and "magnetic" contributions to the total vacuum energy).

4 How topologically nontrivial threads arise inside the discrete non-Abelian vacuum manifold.

The outlines of this topic were us projected in Introduction.

It should be begun from the fact that the Minkowskian Higgs model [9, 10, 11] with vacuum BPS monopoles is the specific model where the duality in specifying the sign of the vacuum "magnetic" field $\mathbf{B}$ takes place. This duality is just set by the Bogonol'nyi equation (1.1) [11].

This duality implies automatically the duality in specifying the signs of magnetic charges $\mathbf{m}$, given by the general formula [9]

$$\mathbf{m} = \frac{1}{4\pi} \int_{\Gamma} d\mathbf{S} \cdot \mathbf{B} = \frac{1}{8\pi} \int d^3x \partial_i \{ \epsilon^{ijk} < F_{jk}^b, \Phi_b > a^{-1} \}. \quad (4.1)$$

Note that latter Eq. is irrelatively correct, actually, to the concrete choice of the vacuum geometry: either "continuous" or "discrete" one, in the quested Minkowskian Higgs model.

Really, due to the Bianchi identity $D B = 0$, the definition (4.1) of magnetic charges $\mathbf{m}$ may be recast formally to the look

$$\mathbf{m} = \frac{1}{4\pi} \int_{\Gamma} d\mathbf{S} \cdot \mathbf{B} \sim \frac{1}{4\pi} \int_V \text{div } \mathbf{B} dV = 0,$$

and this integral does not depend on the choice of the space-like surface in the (Minkowski) space.

However, the actual divergence of the "magnetic" tension $\mathbf{B}$ at the origin of coordinates in all the Minkowskian Higgs models (for instance, [10, 33, 34]) implies that the latter

integral is different from zero as the origin of coordinates lies inside the chosen space-like surface Γ .

Furthermore, the above duality [11] in specifying the sign of the vacuum "magnetic" field $\mathbf{B}$ affects also the generator $h(\Phi)$ [9] of the residual $U(1)$ gauge group in the studied Minkowskian Higgs model.

It is so since $h(\Phi)$ may be read easily from Eq. (4.1).

Now let us return to the simpler case of "continuous" ($\sim S^2$) vacuum geometries in Minkowskian Higgs models.

In this case the appropriate degeneration space: say, R , is one-connected:

$$\pi_0(H) = \pi_1(R) = 0,$$

and one can always choose, in the unique way, a continuous branch of the function $h(\Phi)$ on R .

Another situation is when the degeneration space R is multi-connected ("discrete"), i.e. when

$$\pi_0(H) = \pi_1(R) \neq 0,$$

and therefore there exist thread topological defects in the gauge theory in question (in particular, in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac [4]).

On the one hand (as we have seen this above), the nontrivial isomorphism (2.1) [9] implies the existence of thread topological defects inside the discrete vacuum manifold R .

On the other hand, it turns out (see §Φ13 in [9]) that the duality in the choice of the sign of $h(\phi)$ (and therefore also the duality [11] in the choice of the direction of $\mathbf{B}$) may become unavoidable.

This occurs, in particular, when one can invert the function $h(\Phi)$ via the gauge transformations

$$\tilde{h}(\Phi) = \alpha h(\Phi) \alpha^{-1} = -h(\Phi), \quad (4.2)$$

with

$$\alpha = \exp(2\pi M) \quad (4.3)$$

and the element M belonging to the Lee algebra of the group G_1 of rigid transformations compensating changes in the vacuum (YM-Higgs) thread configuration (Φ^a, A_μ^a) at rotations around the axis z of the chosen reference frame.

The general look of an element $g_\theta \in G_1$ may be set by Eq. (2.5) [9].

Herewith our above discussion about thread topological defects inside degeneration spaces in the liquid helium II models (turning [13] and at rest [15]) prompts us the idea to associate the rectilinear threads (as infinite thin tubes of infinite lengths) with the above axial symmetry.

In the monograph [9] (in §Φ12) the strict proof was given that the unitary group G_1 is a possible symmetry for alike threads (by analogy with the group $O(2)$ of orthogonal 2×2 matrices with determinants ± 1 , one utilizes in the liquid helium II turning model [13]).

Omitting details, note that the proof of this statement comes [9] to the proof that in each topological class, rectilinear threads exist giving a finite contribution to the appropriate energy integrals (due to first-order phase transitions occurred) and satisfying the appropriate equations of motion.

The topological type of a thread defect is determined by the group G_1 : more exactly by its generator M .

To prove this statement, we betake to the argument of the holonomies group [7, 9].

Let us rewrite an element (3.16) of the holonomies group $H \simeq U(1)$ in the shape

$$\alpha = (P \exp(-\oint A_\mu dx^\mu))^{-1}. \quad (4.4)$$

Due to the standard Pontryagin degree of a map theory, elements α of the holonomies group H depend on integers n : $\alpha \equiv \alpha(n)$ ($n \in \mathbf{Z}$).

Going over then to the cylindrical coordinates, we come to Eq. (2.9) [9], implicating rectilinear (topologically nontrivial) threads A_θ , may be represented as (2.4).

Alternatively, A_θ can be represented also as [9]

$$A_\theta(\rho) = M + \beta(\rho), \quad (4.5)$$

where the function $\beta(\rho)$ approaches zero as $\rho \rightarrow \infty$.

This also ensures [9] a finite energies densities in Minkowskian Higgs models involving (topologically nontrivial) threads $A_\theta(\rho)$ and testifies to their location in the region $\rho \rightarrow 0$, i.e. (intimately) near the origin of coordinates.

Thus at the spatial infinity elements $\alpha(n) \in H$ acquire the quested look ¹²

$$\alpha(n) = \exp(2\pi M). \quad (4.6)$$

¹²In Ref. [9] the following general proof of Eq. (4.2) was given.

Firstly, there was assumed that the elements $\alpha(n)$, (4.4), of the holonomies group $H \simeq U(1)$ belong to its adjoint representation.

Secondly, the "generalized case" when $U(1)$ is the subgroup of an invariant subgroup $\tilde{H} = K \otimes U(1)$ in the initial gauge symmetry group G (with K locally isomorphic to a product of simple non-Abelian Lee groups) was discussed in Ref. [9].

A continuous family α_t of elements of $\tilde{H} \subset G$ was considered connecting the element $\alpha(n)$, (4.4), with the unit element of G (it was put $0 \leq t \leq 1$, $\alpha_1 = \alpha$, $\alpha_0 = 1$, $\alpha h(\Phi_0)\alpha^{-1} = -h(\Phi_0)$; with Φ_0 being the fixed point in the vacuum manifold).

At assuming [9] that the elements $\alpha(n)$, (4.4), belong to the adjoint representation of the group G , the generator of the group $U(1) \subset \tilde{H}(T(\alpha_t)\Phi_0)$ (with $T(g)$ being the transformation corresponding to the group element $g \in G$) can be taken as

$$\tilde{H}(T(\alpha_t)\Phi_0) = \alpha_t h(\Phi_0) \alpha_t^{-1}.$$

Of course, another choice is possible, with this generator of the opposite sign, but if the purpose is to have a continuous one-valued branch of the function $h(\Phi)$, the sign of the generator $h(\alpha_t(\Phi_0))$ is no longer arbitrary.

Then at $t = 1$ one has

$$h(\Phi_0) = h(\alpha(\Phi_0)) = \alpha h(\Phi_0) \alpha^{-1} = -h(\Phi_0).$$

In latter Eq. the fact [9] that all the subgroups $\tilde{H}(\Phi)$ in G are mutually conjugate (i.e. that there exist an intrinsic automorphism in G mapping $\tilde{H}(\Phi)$ into a $\tilde{H}(\Phi')$) was taken into account.

We see that the dependence of gauge fields A on the angle θ disappears at the spatial infinity.

It is equivalent to damping, at the spatial infinity, various rotary effects, associated in Minkowskian non-Abelian theories with thread topological defects.

In particular, it is correctly for the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac [4] and involving the "discrete" vacuum geometry (3.9).

On the other hand, the function $\beta(\rho)$ should be different from zero effectively only in the ϵ -neighbourhood of the origin of coordinates to ensure in this infinitesimal region the pointed above collective rotary phenomena in the appropriate Minkowskian YM vacuum via the nontrivial dependence of $A_\theta(\rho, \theta, z)$ on θ .

In the ϵ -neighbourhood of the origin of coordinates (i.e. at $\rho \rightarrow 0$), the function $\beta(\rho)$ would be singular. It is the necessary thing [45] to ensure gauge transformations (4.2) [9], changing the sign of the $U(1)$ generator $h(\Phi)$ (that implies changing the sign of the magnetic charge $\mathbf{m}$: the resulting flip of the magnetic charge $\mathbf{m}$ is $2m$, from $-m$ to m).

Due to the "total" conservation law for the magnetic charge $\mathbf{m}$, YM thread solutions $A_\theta(\rho, \theta, z)$ [9] would bear the "compensating" magnetic charge $-2m$ [45].

The distinctive property of such "compensating" magnetic charges $-2\mathbf{m}$, in comparison with "ordinary" magnetic charges $\mathbf{m}$ (associated with vacuum BPS monopole solutions), is [37] their "discrete" origin, the presence of (topologically nontrivial) threads in the Minkowskian Higgs model quantized by Dirac [4] under consideration. Finally, this is associated with the "discrete" vacuum geometry (3.9) us assumed to justify the Dirac fundamental quantization of the above model.

The "discrete" magnetic charges (let us denote them as $\mathbf{m}_Z$) connected with (topologically nontrivial) threads are not shielded by the condensate of the Higgs field.

More exactly [37], imagining a particle of the magnetic charge me (with e being the unit magnetic charge), the operator

$$\mathfrak{S}_V = \exp(2\pi i \mathbf{M}_V / \mathbf{m}_Z), \quad (4.7)$$

with $\mathbf{M}_V$ being the magnetic charge distributed in the spatial volume V (specefied via the general formula (4.1) [9]), can be constructed as a long range observable.

Note [45] that there is no gauge invariant way to pinpoint when the abov charge transfer has occured. The field configuration (Φ^a, A_μ^a) is $U(1)$ -invariant, but the direction of the arrow on the A_μ^a field line is gauge dependent. One can move the cut by performing a singular gauge transformation (4.2); this alters the apparent time of the charge transfer without actually changing the physics of the process.

Indeed, above described (vacuum) YM thread modes $A_\theta(\rho, \theta, z)$ [9] would be different from vacuum YM BPS monopoles (for instance, in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac).

Thus it becomes obvious that the assumed existence of a continuous one-valued branch for $h(\Phi)$ is inconsistent.

This proof [9] of Eq. (4.2) can be generalized easy for nontrivial topologies $n \neq 0$. To do this, one would replace $\alpha_0 = 1$ (that corresponds to trivial topologies $n = 0$) with an arbitrary α_0 .

It is associated with the obvious divergence as r^{-2} [10] (c.f. (3.17) [36]) of the vacuum "magnetic" field $\mathbf{B}$ at the origin of coordinates in Minkowskian non-Abelian models involving vacuum BPS monopole solutions.

In this case there is impossible to link YM fields $A_\theta(\rho, \theta, z)$, representing topologically nontrivial (rectilinear) threads in the Minkowskian Higgs model, to vacuum BPS monopoles quantized by Dirac, with vacuum YM BPS monopole solutions (with same topological numbers), at least in a smooth wise (that is connected with the $\sim \partial_\mu A_\nu$ items always entering the "magnetic" field $\mathbf{B}$ in YM theories).

On the other hand, the value $\langle B^2 \rangle$, the vacuum expectation value of the "magnetic" field $\mathbf{B}$ squared, can serve [8] as the order parameter in Minkowskian Higgs models with YM fields.

Herewith its nonzero value, $\langle B^2 \rangle \neq 0$, is the sign violating the initial $SU(2)$ gauge symmetry group down to its $U(1)$ subgroup, and it is always the second-order phase transition [6, 28].

If the Minkowskian Higgs model with YM fields implies the continuous $\sim S^2$ vacuum geometry, the possible (as in the BPS or 't Hooft-Polyakov models) divergence of $\mathbf{B}$ at the origin of coordinates testifies [6] in favour of point hedgehog topological defects located just at the origin of coordinates.

The origin of such topological defects is in the nontrivial isomorphism (2.2).

This singularity of (vacuum) "magnetic" fields can be removed by regularising appropriate energy integrals.

Also there are no contributions from point hedgehog topological defects into these energies integrals unlike the case of first-order phase transitions when such contributions appear (as, for instance, ΔE [13], (3.18), in the case of a liquid helium II specimen turning together with the cylindrical vessel where it is contained).

Unlike the former case, gaps in the directions of $\mathbf{B}$ and $\mathbf{B}_1$, (2.6), in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac and involving the "discrete" vacuum geometry (3.9) according to our assumption are evidences in favour of the first-order phase transition occurring in this model.

This is in a good agreement with the general theory of first-order phase transitions (discussed, for instance, in Ref. [28], in §3.1; see also [6]).

The discontinuous behaviour of plots for order parameters is just the sign of first-order phase transitions occurring in physical theories ¹³.

The above discussed gaps do not influence however the "right" of gauge matrices $\alpha(n)$, (2.9) (taking account of the "discrete" vacuum geometry (3.9), involving [topologically nontrivial] threads), to represent the holonomies group H coinciding with the residual gauge symmetry group $U(1)$ in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac [4].

¹³The discussed gaps in the directions of $\mathbf{B}$ and $\mathbf{B}_1$ discovered in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac is a specific manifestation of singularities associated with vortices presenting in that model. The existence of alike singularities in gnon-Abelian gauge theories quantized by Dirac was predicted as far back in Ref. [16] (in the English version of that paper).

This may be explained by the natural isomorphism between exponential multipliers $\alpha(n)$, referring to (topologically nontrivial) threads arising because of the "discrete" vacuum geometry (3.9), and Gribov topological multipliers $v^{(n)}(\mathbf{x})$, referring to superfluid potential motions in the above model.

Note that the dependence of the theory considered on nontrivial topologies n also disappears at the spatial infinity, i.e. in the infrared region of transferred momenta.

It is the next in turn evidence in favour of the infrared topological confinement (in the spirit [1, 23]) occurring in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac.

This "confluence" of different topologies n in the infrared region of the momentum space is closely related with the tendency of domain walls between different topological sectors of the vacuum manifold R_{YM} to become more "thin" as the spatial volume V approaches infinity.

In this limit, the typical width of a domain wall, $\epsilon(V)$, goes, indeed, to zero, as it was discussed above (see (3.14)).

The above discussed examples of liquid helium II at rest [15] and turning together with the cylindrical vessel where it is contained [13] give us pattern of physical systems in which first-order phase transitions occur.

The "centre of gravity" of the enumerated models is in the appearance of (rectilinear) threads generated appropriate vortices, i. e. thread topological defects.

On the other hand, it was demonstrated above (repeating the arguments [13]), in the case of a liquid helium II specimen turning together with the cylindrical vessel where it is contained, the "rotary" contribution ΔE , (3.18), supplements the typical (diagonalized) Bogolubov Hamiltonian $\hat{\mathcal{H}}$ [41].

It is the sign of the first-order phase transition taking place in the liquid helium II turning model [13].

Herewith according to the modern terminology [6], this "rotary" contribution ΔE , (3.18), is the *latent heat*, while the thermodynamic phase in whose framework collective solid potential rotations [13, 15] in a liquid helium II specimen turning together with the cylindrical vessel where it is contained coexist with superfluid motions setting by the Hamiltonian $\hat{\mathcal{H}}$ is characterized by the *supercooling* phenomenon.

In this case superfluid motions inside a liquid helium II specimen represent the *stable state* corresponding to the *second-order* phase transition occurred therein beforehand and coming to violating the $U(1)$ gauge symmetry of the helium Hamiltonian $\hat{\mathcal{H}}$ [41]¹⁴ (in the

¹⁴The transparent evidence in favour of the second-order phase transition occurring in a helium: actually, it is the transition between the liquid helium I high-temperature and liquid helium II modifications (the latter one possesses manifest superfluid properties described in the Landau-Bogolubov theory [14, 42]) is the discontinuity [46] in the plot of the helium heat capacity in the Curie point $T_c \rightarrow 0$.

For instance [41], in the case of superfluid helium films, the experimental data result

$$C_p(T) \equiv T \left(\frac{\partial S}{\partial T} \right) \approx c_1 + c_2 \exp[c_3/|T - T_c|^{1/2}]$$

for the helium heat capacity $C_p(T)$, with S being the appropriate entropy; c_i ($i = 1, 2, 3$) are constants.

(3.22) wise in the liquid helium II turning case [13]).

In turn, collective solid potential rotations in this specimen are referred to the *metastable state* (in the terminology [28]).

There are concrete computations, given in Ref. [41] with the example of a rested liquid helium II specimen, demonstrating discontinuities in the plot of the vacuum expectation value $\langle |\Xi|^2(t, \mathbf{r}) \rangle$ for the helium Bose condensate wave function Ξ , (1.4), near locations of vortices cores.

Herewith the value $\langle |\Xi|^2(t, \mathbf{r}) \rangle$ can serve quite naturally as the order parameter in the Landau-Bogolubov helium theory [14, 42].

So $\langle |\Xi|^2(t, \mathbf{r}) \rangle = 0$ for the helium I modification [46], corresponding to the initial $U(1)$ gauge symmetry of the helium Hamiltonian $\hat{\mathcal{H}}$ [41], while $\langle |\Xi|^2(t, \mathbf{r}) \rangle \neq 0$ upon this symmetry is violated in the (3.21) wise, that corresponds to the helium II modification.

Then, utilizing Eq. (1.4) [6, 13], the correlation (Green) function

$$G(r) \equiv \langle \Xi(t, \mathbf{r}), \Xi^*(t, 0) \rangle$$

(it is different from zero for the liquid helium II modification) can be recast to the look [41]

$$\begin{aligned} G(r) &= \langle \Xi(t, \mathbf{r}), \Xi^*(t, 0) \rangle \\ &= \Xi_0^2 \exp \left[-\frac{1}{2} \langle [\Phi(t, \mathbf{r}) - \Phi(t, 0)]^2 \rangle \right], \end{aligned} \quad (4.8)$$

where the last line is a general property of Gaussian fluctuations.

To a first approximation, one can neglect fluctuations in the amplitude of Ξ , setting it to be a Ξ_0 . Then the vacuum expectation value $\langle |\Xi|^2(t, \mathbf{r}) \rangle$ of the helium Bose condensate wave function Ξ will equal to $G(0) = (\Xi_0)^2$.

On the other hand, the above discussed expression [15] for topological indices $n \in \mathbf{Z}$ via circular velocities $\mathbf{v}^{(n)}$ is mathematically equivalent to [41]

$$\frac{m}{2\pi\hbar} \oint_{\Gamma} \nabla \Phi \, d\mathbf{l} = \int_{\Omega} d^2r \, n_v(\mathbf{r}), \quad (4.9)$$

with the vortex "charge density" given as

$$n_v(\mathbf{r}) = \sum_{\alpha=1}^N n_{\alpha} \delta(\mathbf{r} - \mathbf{r}_{\alpha}) \quad (4.10)$$

for a collection N of vortices located at positions $\{\mathbf{r}_{\alpha}\}$ with integer charges $\{n_{\alpha}\}$. Without loss of generality, it may be set $z = 0$ in all the calculations [41] following. In (4.9), Eq. (1.3) [6, 13], with appropriate replacing $\mathbf{v}_0 \rightarrow \mathbf{v}^{(n)}$, was utilized.

Applying then the Stokes formula to (4.9), one find upon integrating:

$$\begin{aligned} \epsilon_{ij} \partial_i \partial_j \Phi(t, \mathbf{r}) &= \partial_x \partial_y \Phi - \partial_y \partial_x \Phi \\ &\approx n_v(\mathbf{r}). \end{aligned} \quad (4.11)$$

To cast this Eq. in a more familiar form, one can introduce [41] the value $\tilde{\Phi}(t, \mathbf{r})$ dual to $\Phi(t, \mathbf{r})$:

$$\partial_i \Phi(t, \mathbf{r}) = \epsilon_{ij} \partial_j \tilde{\Phi}(t, \mathbf{r}). \quad (4.12)$$

Then

$$\nabla^2 \tilde{\Phi}(t, \mathbf{r}) = n_v(\mathbf{r}). \quad (4.13)$$

In particular,

$$\nabla^2 \tilde{\Phi}(t, \mathbf{r}) = 0 \quad (4.14)$$

in the case when vortices are absent, i.e. in the case of purely superfluid motions inside a liquid helium II specimen.

Eq. (4.13) is the particular case of the Poisson equation. Thus it permits its

$$\tilde{\Phi}(t, \mathbf{r}) = \sum_{\alpha} n_{\alpha} G(\mathbf{r}, \mathbf{r}_{\alpha}) \quad (4.15)$$

solution, where the Green function $G(\mathbf{r}, \mathbf{r}_{\alpha})$ (defined as on the left-hand side of Eq. (4.8)) satisfies

$$\nabla^2 G(\mathbf{r}, \mathbf{r}_{\alpha}) = \delta(\mathbf{r} - \mathbf{r}_{\alpha}). \quad (4.16)$$

For $|\mathbf{r} - \mathbf{r}_{\alpha}|$ large enough and both the points far from any boundaries, the Green function $G(\mathbf{r}, \mathbf{r}_{\alpha})$ has the look (see §11.8 in [47])

$$G(\mathbf{r}, \mathbf{r}_j) \approx \frac{1}{2\pi} \ln \left(\frac{|\mathbf{r} - \mathbf{r}_j|}{\xi_0} \right) + C \quad (4.17)$$

on the $z = 0$ plane, where C is a constant which contributing to the vortex core energy.

The origin of the parameter ξ_0 entering Eq. (4.17) is following [41]. It is the characteristic length related to the coefficients of the first two terms of Eq. [13]

$$\frac{F}{T} = \int d^2 r \left[\frac{1}{2} A |\nabla \Xi|^2 + \frac{1}{2} a |\Xi|^2 + b |\Xi|^4 + \dots \right] \quad (4.18)$$

for the helium free energy F .

In this case ξ_0 can be defined as [41]

$$\xi_0 = \sqrt{A/|a|}, \quad (4.19)$$

with a given as $a = a'(T - T_c)$.

As it follows from (4.17), at setting $C = 0$, the Green function $G(\mathbf{r}, \mathbf{r}_j)$ diverges logarithmically when $|\mathbf{r} - \mathbf{r}_j| \rightarrow \infty$, while it approaches zero when

$$|\mathbf{r} - \mathbf{r}_j| \rightarrow \xi_0.$$

Additionally, $G(\mathbf{r}, \mathbf{r}_j)$ diverges logarithmically also at $|\mathbf{r} - \mathbf{r}_j| \rightarrow 0$. The latter property of $G(\mathbf{r}, \mathbf{r}_j)$, as we shall see soon, is very important.

The just performed analysis of $G(\mathbf{r}, \mathbf{r}_j)$ allows to draw the series important conclusions about the phase $\tilde{\Phi}(t, \mathbf{r})$ of the helium wave function Ξ (taking over the complete collection N of topologies: including the trivial one $s_\alpha = 0$, corresponding to superfluid motions [14, 15], in a liquid helium II specimen at rest).

So, due to (4.8),

$$\langle |\Xi|^2(t, \mathbf{r}) \rangle \equiv \langle \Xi(t, \mathbf{r}), \Xi^*(t, \mathbf{r}) \rangle \approx G(0) = 1/N \sum_{\alpha} \langle |\Xi^2|_{\alpha}(t, \mathbf{r}) \rangle, \quad (4.20)$$

serving the order parameter in the helium theory, diverges logarithmically at each index $\alpha(n)$ ($n \in \mathbf{Z}$) in the same (4.17) [41] sense that the Green function $G(\mathbf{r}_\alpha, \mathbf{r}_\alpha) \approx G(0)$, i.e. in the points $\{\mathbf{r}_\alpha\}$ where quantum vortices are located in a liquid helium II specimen at rest ¹⁵.

This is associated immediately with (rectilinear) quantum vortices arising spontaneously [15] in a liquid helium II specimen at rest. Herewith also the phase $\tilde{\Phi}(t, \mathbf{r})$ of the helium wave function Ξ possesses the same behaviour that the Green function $G(\mathbf{r}, \mathbf{r}_j)$ (it is correctly due to (4.15)).

The said is true for nontrivial vortices topologies $n \neq 0$ due to (4.15). However for $n = 0$, i.e. in the "superfluid case", $\tilde{\Phi}(t, \mathbf{r})$ becomes an uncertain value when $|\mathbf{r} - \mathbf{r}_j| \rightarrow \infty$ (or when $|\mathbf{r} - \mathbf{r}_j| \rightarrow 0$).

On the other hand, the phase $\tilde{\Phi}(t, \mathbf{r})$ (or $\Phi(t, \mathbf{r})$) of the helium wave function Ξ , (1.4), disappears from Eq. (4.20).

Summarizing up, now we can assert that in the presence of (rectilinear) quantum vortices [15] in a liquid helium II specimen, the order parameter $\langle |\Xi|^2(t, \mathbf{r}) \rangle$ in the helium model suffers discontinuities of the logarithmic nature in the points $\{\mathbf{r}_\alpha\}$ in which these (rectilinear) quantum vortices are located (the long distances logarithmic singularities of $G(0)$ are not associated, probably, with quantum vortices).

The discovered discontinuity (of the logarithmic nature) in the plot of the vacuum expectation value $\langle |\Xi|^2(t, \mathbf{r}) \rangle$ for the helium Bose condensate wave function Ξ , serving the order parameter in the helium theory, just testifies in favour of the first-order phase transition occurring in a liquid helium II rested specimen with arising therein quantum vortices.

There are, indeed, lot of distinctions between the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac [4] and the liquid helium II at rest theory [15] (the same concerns also the liquid helium II turning model [13], involving rigid $O(2)$ rotations).

But, in spite of these distinctions, the discontinuities in the appropriate order parameters: $\langle |\Xi|^2(t, \mathbf{r}) \rangle$ in the helium theory and $\langle B^2 \rangle$ in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac, can be considered as the common trait

¹⁵In (4.20) the Green function $G(0)$ is different from that given via Eq. (4.8) and involving the constant amplitude Ξ_0 . Now the amplitude of $\Xi(t, \mathbf{r})$ wouldn't be constant, and this implies that $G(0)$ has the nontrivial look $G(\mathbf{r}_\alpha, \mathbf{r}_\alpha)$ [41], (4.17).

of the both models associated with rotary effects (including the "discrete" vacuum geometry and thread topological defects).

Moreover, drawing a parallel between the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac and the liquid helium II at rest theory [15], one can refer collective solid rotations inside the appropriate BPS monopole vacuum to the metastable thermodynamic phase, characterised by the latent heat equal to the "electric" energy $\sim E^2$. This "electric" energy is described correctly by the free rotator action functional (1.11).

Simultaneously, superfluid potential motions inside this vacuum, involving the "magnetic" energy $\sim B^2$, may be referred to the stable thermodynamic phase.

It is again the supercooling situation.

As it was discussed in Section 2, on the face of it, there is a contradiction between the rotary effects in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac (these effects, including collective vacuum rotations, are determined by the action functional (1.11)) and the manifest potential nature of the Minkowskian BPS monopole vacuum.

More exactly, this potential nature of the Minkowskian BPS monopole vacuum (suffered the Dirac fundamental quantization [4]) comes, in particular, to the colinearity (2.8) [16] of vectors $\mathbf{B}$ and $\mathbf{E}$ (vacuum "magnetic" and "electric" fields, respectively).

This is associated with the explicit look (1.13), (1.15) [5, 7, 8, 12] for vacuum "electric" monopoles $\mathbf{E}$, can be expressed through vacuum Higgs BPS monopole modes $\Phi_{(0)}(\mathbf{x})$.

Also the vacuum "magnetic" field can be expressed through vacuum Higgs BPS monopole modes $\Phi_{(0)}(\mathbf{x})$, now through the Bogomol'nyi equation (1.1).

There is however a way to solve the above contradiction.

It turns out that, side by side with (topologically nontrivial) threads A_θ , one can construct also specific vacuum Higgs z -invariant solutions [9], having the look (2.7) in the cylindrical coordinates, such that the vacuum Higgs field $\Phi_a^{(n)}$ specified by (2.7) merges with the appropriate Higgs vacuum BPS monopole solution in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac [4], herewith belonging to the same topology n .

Additionally, we should also claim that the covariant derivative $D\Phi$ of any vacuum Higgs field $\Phi_a^{(n)}$ specified via (2.7) merges with the covariant derivative of such a vacuum Higgs BPS monopole solution.

In other words, vacuum Higgs solutions (2.7), corresponding to the above described axial symmetry G_1 , would merge with appropriate Higgs vacuum BPS monopole solutions actually in a smooth wise.

All this can be achieved by the appropriate choice of the function $\phi(\rho)$ in (2.7) (the latter one satisfies the spatial asymptotic (2.7) [9]).

By this the goal is pursued to link $D\Phi_a^{(0)}$ to vacuum "electric" monopoles (1.13) in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac ¹⁶.

¹⁶As it was explained above, the plot of $D\Phi_a^{(0)}$ in the thread topological defects region inside the

Indeed, this confluence of vacuum Higgs solutions would occur already at the notably lesser scale of distances: at the hadronic distances scales, ~ 1 fm., to ensure the correct infrared behaviour [23] of quark Green functions in the Minkowskian (constraint-shell) QCD.

Actually, the following picture takes place.

As it was noted in Ref. [20], Higgs BPS monopole solutions disappear at the origin of coordinates (in its infinitesimal neighbourhood of the effective radius ϵ).

In this situation the continuous and smooth prolongation of Higgs vacuum BPS monopole solutions with appropriate Higgs modes $\Phi_a^{(n)}$ [9], (2.7), seems to be quite natural. It is controlled by the proper choice of $\phi(\rho)$.

It is just the way solving the above discussed problem how to co-ordinate the "colinearity condition" (2.8) [16] (reflecting the superfluid nature of the Minkowskian BPS monopole vacuum suffered the Dirac fundamental quantization [4]) with collective solid rotations inside this vacuum.

Really, Higgs (vacuum) modes $\Phi_a^{(n)}$, (2.7), refer to the *rigid* axial symmetries group G_1 [9].

This group, in turn, compensates *local* rotations of the vacuum (YM-Higgs) configuration $(\Phi_a^{(n)}, A_\mu^a)$ around the axis z (just such rotations are apprehended as rectilinear vortices).

On the other hand, this vacuum (YM-Higgs) configuration can be linked, in a continuous and smooth wise in its "Higgs" part, to Higgs vacuum BPS monopoles, disappearing at the origin of coordinates [20].

By that two things are ensured.

Firstly, the continuity (and smoothness) achieved now for Higgs vacuum solutions, "legalizes" collective solid rotations inside the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac [4] (as a sign of thread topological defects inside the appropriate vacuum manifold R_{YM}) over the infinitely narrow cylinder of the effective diameter ϵ along the axis z . These rotations are described correctly by the action functional (1.11) [5].

Secondly, simultaneously, the Gauss law constraint (1.8) and the "colinearity condition" (2.8) are satisfied outward of this infinitesimal region with smooth vacuum "electric" monopoles (1.13).

On the other hand, being a continuous and smooth value now, $\langle \Phi^{(0)} \rangle^2$ (with all its Gribov copies $\langle \Phi^{(n)} \rangle^2$; see e.g. [2]) cannot be considered as an order parameter in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac since alike order parameter would suffer singularities in coordinate points where vortices (thread topological defects) are located inside the vacuum manifold R_{YM} (cf. our above discussion [41] about quantum vortices in a liquid helium II specimen).

vacuum manifold R_{YM} does not coincide with the plot of the vacuum "magnetic" field $\mathbf{B}$, given by the Bogomol'nyi equation (1.1), i.e. via vacuum Higgs BPS monopole solutions. This vacuum "magnetic" field $\mathbf{B}$ diverges as $1/r^2$ [10] at the origin of coordinates.

Thus $\langle B^2 \rangle \sim \langle D \Phi^{(n)} \rangle^2$ [9] rather than $\langle \Phi^{(n)} \rangle^2$ serves as an order parameter in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac. It is the specific of the first-order phase transition occurring in that model and involving the discrete vacuum geometry (3.9). This distinguishes it from the Minkowskian Higgs models involving the continuous $\sim S^2$ vacuum geometry, where just the “Higgs” vacuum expectation value $\langle \Phi \rangle^2$ serves as the order parameter.

Note that also in the “classical” Minkowskian (YM-Higgs) model with vacuum BPS monopoles [6, 9, 10, 11], involving the “continuous” $\sim S^2$ vacuum geometry and in which only the second-order phase transition occurs, $\langle B^2 \rangle$ can serve as an order parameter, but now side by side with $\langle \Phi^2 \rangle$.

When we view very attentively Eq. (1.12) [5] for the “rotary momentum” I of the BPS monopole vacuum, we can note that the integration there is carried out over the infinite spatial volume $V \rightarrow \infty$.

At neglecting the infinitesimally small interval $[0, \epsilon(\infty)]$ in this integral, one get nevertheless the same result.

More exactly, the integral (1.12) may be evaluated in the following way [5]:

$$I \simeq \frac{4\pi^2}{\alpha_s} \int_{\epsilon}^R dr \frac{d}{dr} \left(r^2 \frac{d}{dr} f_0^{BPS}(r) \right); \quad R \rightarrow \infty \quad (4.21)$$

It involves the “Higgs” BPS ansatz [7, 8]

$$f_0^{BPS}(r) = \left[\frac{1}{\epsilon \tanh(r/\epsilon)} - \frac{1}{r} \right],$$

disappearing [20] at the origin of coordinates.

We see that the extrapolation of Higgs vacuum BPS monopoles inward the interval $[0, \epsilon]$, where the potential superfluid nature of the Minkowskian BPS monopole vacuum suffered the Dirac fundamental quantization [4] is violated, does not affects the integrals (1.12), (4.21) so long as the integration limits in Eq. (4.21) are $[\epsilon, \infty]$.

This is just the region where the potential superfluid nature of that vacuum takes place.

It is determined, in the Dirac fundamental quantization scheme [4], by the Bogomol’nyi equation (1.1), YM Gauss law constraint (1.8) and Gribov ambiguity equation (1.2) as the logical consequence [2, 7, 8] of the Bogomol’nyi equation (1.1).

As to the interval $[0, \epsilon]$, integrating inside this interval comes to integrating over the base of the infinitely narrow cylinder along the axis z (in the chosen rest reference frame), so long as the Higgs BPS ansatz $f_0^{BPS}(r)$ is a function of the distance r only. Thus the interval $[0, \epsilon]$ would result a vestigial contribution to the “rotary momentum” I , (1.12), and one can neglect it actually.

On hand thus a delimitation between superfluid potential motions and collective solid rotations inside the Minkowskian BPS monopole vacuum suffered the Dirac fundamental quantization [4].

Superfluid potential motions therein (controlled by the Bogomol'nyi and Gribov ambiguity equations) are referred actually to the spatial interval $r \in [\epsilon(\infty); \infty]$, while collective solid rotations to the spatial interval $r \in [0, \epsilon(\infty)]$.

This delimitation is the next in turn evidence in favour of the first-order phase transition occurring in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac and coming to the coexistence of two thermodynamic phases inside the Minkowskian BPS monopole vacuum suffered the Dirac fundamental quantization. The thermodynamic phases of superfluid potential motions and collective solid rotations are just such phases.

To this stage of investigations about the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac, we cannot conclude regarding the concrete nature of the boundary between the mentioned thermodynamic phases as well as regarding various surface effects.

At the same time, we should like note that the transparent way to verify that the first-order phase transition occurs in any physical model is to compute the appropriate surface tension integral [13]

$$\alpha_{12} = \int_{-\infty}^{\infty} (f_1 - f_2) dx, \quad (4.22)$$

where f_1 and f_2 are the free energies densities referring to two thermodynamic phases coexisting at the first-order phase transition occurring in the given physical model.

Meanwhile, already now some conclusions may be drawn concerning to the domain structure of the Minkowskian vacuum manifold R_{YM} , (3.9). It is just associated with the $\epsilon(V) \sim V^{-1}$ dependence [7, 8].

So intimately to the axis z of the chosen rest reference frame, domain walls become very wide (the typical wide of a domain wall ibid of the $\sim \epsilon(0) \rightarrow \infty$ order, while alone topological domains become microscopically thin.

One can assume herewith that in the mentioned spatial region, topological domains inside R_{YM} (at neglecting their thickness) possess approximately the geometry of planes (perpendicular to the axis z). This implies also concentrating rectilinear threads A_θ inside these highly thin topological domains.

Vice verse, at the spatial infinity topological domains merge (as it was noted above), while walls between them become infinitely thin (of the typical wide $\epsilon(\infty) \rightarrow 0$) and this promotes the infrared topological confinement [23] of "large" multipliers $v^{(n)}(\mathbf{x})$ in the (quark, gluonic) Green functions in all the orders of the perturbation theory.

Actually, since the confluence of topological domains would occurs already at the hadronic distances scales, ~ 1 fm., one can speak that only in the spatial region intimately near the axis z of the chosen rest reference frame the "discrete" vacuum geometry R_{YM} , (3.9), in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac [4] is different from the continuous $\sim S^2$ vacuum geometry, (3.1), proper to the "classical" Minkowskian Higgs model with BPS monopoles [9, 10, 11], involving the "heuristic" FP quantization scheme [35], associated with fixing the $A_0 = 0$ gauge [6].

Nevertheless, this distinction in vacuum geometries plays the crucial role, as we see from the current discussion, for understanding the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac.

The important consequence of the presence of topologically nontrivial threads in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac [4] is annihilating [9] two equal magnetic charges $\mathbf{m}_1 = \mathbf{m}_2 = \mathbf{m}(n) \neq 0$ ($n \in \mathbf{Z}$) colliding at crossing a rectilinear topologically nontrivial thread $A_\theta(n)$: (2.4), (4.5).

This effect is the particular case of changes in the relative sign of two magnetic charges as the integration region Γ in (4.1) intersects a (rectilinear) thread.

Suppose that the magnetic charge $|\mathbf{m}| = m_1$ is concentrated to the left of the thread $A_\theta(n)$, while the magnetic charge $|\mathbf{m}| = m_2$ is concentrated to the right of this thread. Herewith we concentrate these magnetic charges in points.

Concentrating magnetic charges in points does not change the matter, despite magnetic charges are distributed volume actually due to (4.1).

Due to (4.2), the both magnetic charges are specified to within their signs.

One can draw a closed curve connected the both magnetic charges (see Fig. 15 in [9]). It consists of two lines, γ_1 and γ_2 , between the charges, one rounds in the opposite directions.

This implies that the vacuum "magnetic" tension $\mathbf{B}$ (specifying via the Bogomol'nyi equation (1.1)) changes its sign depending on the "direction": either $\vec{\gamma}_1$ or $\vec{\gamma}_2$, in which one crosses the thread.

Really, we should always consider the relative sign of the charges $\mathbf{m}_1$ and $\mathbf{m}_2$: more precisely, our interest is the difference $m_1 - m_2$ as we move in the "direction" $\vec{\gamma}_1$; contrariwise, our interest is the difference $m_2 - m_1$ as we move in the opposite "direction" $\vec{\gamma}_2$.

Taking account of the duality (4.2) in the definition of a magnetic charge, we encounter few "alternatives" for the values of the relative magnetic charge in both the cases.

The most interesting case is when $m_1 = m_2$. In this case some two vacuum BPS monopoles belonging to the one fixed topological class $n \in \mathbf{Z}$ annihilated mutually at approaching each other crossing the topologically nontrivial thread A_θ lying in the Minkowskian YM vacuum manifold R_{YM} , (3.9) (the same is correctly also for excitations over the BPS monopole vacuum).

The above reasoning demonstrates that magnetic charges are not maintained in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac.

Magnetic charges cease then to be motion integrals; at the same time they are still topological invariants: continuous deformations do not change the difference $m_2 - m_1$.

In the monograph [9], in §Φ13, there was proposed the following, more general, definition of a magnetic charge, taking account of rectilinear (topologically nontrivial) threads in appropriate non-Abelian gauge models.

One considers a membrane pulling on the rectilinear thread (now treated as a geometrical object in the manner us discussed above) lying along the axis z of the chosen

coordinate system. Then one removes this membrane from the Minkowskian space. This results a one-connected set, on which one can choose a fixed continuous branch of the magnetic tension $\mathbf{B}$.

In fact, this "trick" comes to the restriction of the integration region in Eq. (4.1) for a magnetic charge by the "left" (respectively, "right") semi-space of the Minkowskian space with respect to the axis z . This ensures that magnetic charges are maintained in each of these semi-spaces taken separately.

In the present study we don't have for our object deriving the concrete contribution from the above discussed annihilating effect [9] for equal topologically nontrivial magnetic charges colliding at crossing a thread A_θ in the total action functional (Hamiltonian) of the quested Minkowskian Higgs model with vacuum BPS monopoles, although such contribution exists undoubtedly.

Nevertheless, now we should like make some remarks may be helpful for finding such contribution.

For instance, in the paper [48] possible interactions between (YM) magnetic monopoles and antimonopoles were studied involving forming so-called *monopole molecules*.

Herewith only two types of monopole molecules are possible [48], reflecting ordinary Feynman rules for gauge non-Abelian fields (see e.g. §9.1 in [24]).

There are, firstly, confined colourless magnetic monopoles M and $\tilde{M}$ (with the symbol M referring to a monopole with the magnetic charge $\mathbf{m}$ and the symbol $\tilde{M}$ referring to an antimonopole with the magnetic charge $-\mathbf{m}$, respectively). These monopole molecules were also referred to as monopole-antimonopole *mesons* in Ref. [48]. The well-known monopole-antimonopole confinement and annihilation mechanism via Abrikosov threads (strings) of the "magnetic" field [9, 36, 49] is implicated here (see also §6.2 in [28]).

On the phenomenological level, such Abrikosov threads [9, 36, 49] can be describes with the aid of the *Mandelstam* linearly increasing potential $E \sim Kr$ [24] (with K being the tension of the Abrikosov string).

Secondly [48, 24], confined colour triplet and antitriplet of monopoles are also possible in the nature. Herewith MMM (respectively, $\tilde{M}\tilde{M}\tilde{M}$) states were referred to as *baryons* in [48].

The presence of topologically nontrivial threads in a gauge non-Abelian theory (for instance, rectilinear threads (2.4) [9]), implying the annihilation of monopole pairs with *equal* magnetic charges colliding at crossing such a (rectilinear) topologically nontrivial thread, changes substantially the matter in comparison with the case of the Abrikosov mechanism [28, 36, 48] of the monopole-antimonopole confinement.

The former case is rather similar to the case of confined colour triplet and antitriplet of monopoles (where the role of the one field in such triplets is played by the rectilinear thread A_θ). The only important specific is that the rectilinear thread A_θ would bear always the magnetic charge $\mathbf{m}_Z = -2\mathbf{m}$ [45, 37] due to (above discussed) reasoning maintaining the magnetic charge.

Note that this "baryonic" way annihilating equal magnetic charges colliding with the rectilinear thread A_θ is somewhat alternative to one discussed in Ref. [9]. In the latter

case the $\mathbf{m}$ and $-\mathbf{m}$ pairs forming upon crossing by the one of the magnetic charges the thread A_θ annihilate rather in the “mesonic” $\tilde{M}M$ way as it was described in [48].

The serious objection against this “mesonic” annihilation way is its temperature dependence [28, 48]. More precisely [48], the annihilation of monopole-antimonopole pairs connected with the Mandelstam linearly increasing potential $E \sim Kr$ comes to rotating monopoles about each other inside the bag of the typical size $(g^2T/\hbar^2c^2)^{-1}$ with the velocity exceeding the *thermal velocity* $\sim \sqrt{T/m_M}$, with m_M being the mass of a magnetic monopole. Then during the time m_M/T^2 magnetic monopoles lose all their energy and annihilate.

And moreover, if a $\tilde{M}M$ pair is immersed in the hot YM plasma, this pair is confined by Abrikosov threads of the typical length $\Delta l \sim (e^2T)^{-1}$. Whence, $E \sim K(e^2T)^{-1}$ [28].

One can see thus that the annihilation of monopole-antimonopole pairs connected with the Mandelstam linearly increasing and temperature depending potential $E(T)$ proceeds more quickly at higher temperatures (when lengths of Abrikosov threads decrease) while slower (and perhaps, is suppressed at all) at lower temperatures (in particular, at $T \rightarrow 0$; in particular, the Minkowskian Higgs model with vacuum BPS monopole solutions quantized by Dirac, the topic of the present study, is us considered at such low temperatures).

On the other hand, the computations about the annihilation of monopole pairs with equal magnetic charges (at least at the level of Feynman diagrams) generalizes somewhat the computations [50, 51] about electron-positron pairs in QED.

Herewith unlike QED, where the creation-annihilation processes are forbidden with an only one photon taking part in such processes (by the reason of maintaining the four-momentum in a Feynman diagram involving two fermions and one photon on-shell: see e.g. §25.1 in [51]), in an YM model involving only massless gauge fields, processes with three YM fields are quite permissible.

In particular, it is correctly for the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac, where YM fields, transverse topological Dirac variables A_i^D ($i = 1, 2$), are specified [2] over the light cone $p^2 = 0$ due to the removal (3.11) [5] of temporal YM components.

Note on the way that us discussed [9] colliding processes with magnetic charges $\mathbf{m}$ and (topologically nontrivial) threads in the Minkowskian non-Abelian Higgs model (with vacuum BPS monopoles) quantized by Dirac can be treated as a particular case of the well-known *Aharonov-Bohm effect*.

This effect, pointing to the nontrivial topological structure of the vacuum in a gauge model, was discovered originally in the paper [52]; its good description was given in Ref. [36], in §3.4.

The insurmountable duality [9] in specifying the sign of the vacuum “magnetic” field $\mathbf{B}$ in the Bogomol’nyi equation (1.1) taking place at assuming the discrete geometry (3.9) for the vacuum manifold R_{YM} in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac was us discussed above.

Remember herewith that the crucial point for this “insurmountable duality” is the existence of gauge transformations (2.10) [9] (depending manifestly on thread solutions

A_θ via Eq. (2.9)).

Besides this duality, there is also the manifest symmetry of the "electric" action functional (1.11), squared by the topological momentum P_N (given via Eq. (1.14)), with respect to the changes in the sign of $P_N(k)$ ($k \in \mathbf{Z}$).

This becomes more obvious at recasting (1.11) to the look

$$W_N = \int dt \frac{P_N^2(t)}{2I}. \quad (4.23)$$

Actually the said is equivalent to identifying each two topologies with "opposite signs": k and $-k$. This is again the merit of gauge transformations (4.2).

Thus, with taking account of the "discrete" factorisation (3.3) for the residual $U(1)$ gauge symmetry, the "total" residual gauge symmetry group of the free rotator action (1.11) amounts

$$U_1 = U_0 \otimes \mathbf{Z}/\mathbb{Z}_2. \quad (4.24)$$

The same conclusion about identifying each two topologies with "opposite signs" can be drawn about the vacuum "magnetic" energy [5]

$$\frac{1}{2} \int_\epsilon^\infty d^3x [B_i^a(\Phi_k)]^2 \equiv \frac{1}{2} V < B^2 > = \frac{1}{2\alpha_s} \int_\epsilon^\infty \frac{dr}{r^2} \sim \frac{1}{2} \frac{1}{\alpha_s \epsilon} = 2\pi \frac{gm}{g^2 \sqrt{\lambda}} = \frac{2\pi}{g^2 \epsilon}, \quad (4.25)$$

squared by the vacuum "magnetic" tension $\mathbf{B}$, set, in turn, by the Bogomol'nyi equation (1.1) and responsible for the superfluid properties of the BPS monopole vacuum (suffered the Dirac fundamental quantization [4]). On the other hand, the Bogomol'nyi equation (1.1) [11], associated the vacuum "magnetic" field $\mathbf{B}$ to the Higgs BPS monopole isomultiplet Φ^a , specifies, indeed to within a sign, this vacuum "magnetic" field $\mathbf{B}$ and thus involves the invariance of the vacuum "magnetic" energy (4.25) with respect to changes in signs of magnetic, that is equivalent of *topological*, charges.

The said implies the modification of the "discrete" geometry (3.9) of the vacuum manifold R_{YM} .

Now it may be written down as

$$R'_{YM} = (\mathbf{Z}/\mathbb{Z}_2) \otimes G_0/U_0, \quad (4.26)$$

while the "modified" residual gauge symmetry group in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac is U_1 , given by Eq. (4.24).

This implies also the "modified" discrete representation

$$G_M = G_0 \otimes \mathbf{Z}/\mathbb{Z}_2 \quad (4.27)$$

for the initial gauge symmetry group in this model.

It is easy to see that the topological content of the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac does not change at the modifications (4.24), (4.26), (4.27).

Thus all the kinds of topological defects: thread and point hedgehog ones and domain walls between different topological sectors maintain in that model in spite of the modifications (4.24), (4.26), (4.27).

5 Specific of Higgs phase in Dirac quantisation scheme.

The just discussed annihilation mechanism [9] for (equal) magnetic charges in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac [4] can cause disappearing, in a definite time interval τ , topologically nontrivial magnetic charges $\mathbf{m} \neq 0$ ¹⁷.

Let us suppose that this has occurred indeed. In this case, as it is well known, the Dirac quantization [21, 36, 24] of magnetic and electric charges in the (YM-Higgs) theory, taking the general look [22]

$$e_1 \mathbf{m}_2 + e_2 \mathbf{m}_1 = 2\pi n; \quad n \in \mathbf{Z}; \quad (5.1)$$

for two interacting (BPS, 't Hooft-Polyakov and so on) monopoles in that theory¹⁸, implies for Higgs monopoles to have arbitrary electric charges e associated with the topological number $n = 0$ (and thus with magnetic charges $\mathbf{m} = 0$).

In the review [22] (and in other modern physical literature) such situation in a $SU(n)$ non-Abelian gauge theory (involving the spontaneous breakdown of the initial $SU(n)$ gauge symmetry down to its $U(1)^{n-1}$ subgroup) when gauge fields A possess zero magnetic charges, that implies, in turn, arbitrary electric charges for Higgs modes Φ due to the Dirac quantization [21] of electric and magnetic charges, was referred to as the so-called *Higgs phase*¹⁹.

In the notion "the Higgs phase", one implies however (for instance, in Ref. [22]) that all the magnetic charges are confined by (narrow) Meissner flux tubes, similar to ones in a superconductor [13, 24].

¹⁷Indeed, we must be careful with this assertion because the possibility for (topologically nontrivial) YM modes to penetrate into any topological domain inside the vacuum manifold R'_{YM} .

Such penetration can occur through the spatial region $|\mathbf{x}| \rightarrow \infty$, where domain walls inside R'_{YM} become infinitely thin (as it was discussed above).

Note incidentally that this penetrating YM modes (including vacuum BPS monopole modes) from a one topological domain to another is in a good agreement with the effect [9] raising/increasing the topological charge onto $2k$ ($k \in \mathbf{Z}$) in the presence of (topologically nontrivial) threads inside the vacuum manifold R'_{YM} .

All this complicates somewhat the situation [9] with the annihilation of identical magnetic charges $\mathbf{m}(n)$ in the presence of (topologically nontrivial) threads inside R'_{YM} , implicating additional computations (for instance, it should be confirmed by such computations that the mutual annihilation of equal magnetic charges occurs with the account of the "redundant" YM modes penetrating through the region, at $|\mathbf{x}| \rightarrow \infty$, of confluence different topological domains inside R'_{YM} ; these computations are, perhaps, enough intricate).

¹⁸In the case of a general $SU(N)$ gauge group, one should write down [22]

$$\sum_{i=1}^{N-1} e_i \mathbf{m}_i = 2\pi n$$

instead of Eq. (5.1).

¹⁹Actually [22], these electric charges are screened by the, in average, electrically neutral Higgs Bose condensate in the Higgs phase. The same is correct also for the Minkowskian Higgs model with vacuum BPS monopole solutions quantized by Dirac, where (it can be argued) BPS ansatzes [9, 10] play the role of electric form-factors, that can provide the neutrality of the appropriate Higgs Bose condensate.

In turn, this involves the linearly increasing "Mandelstam" potential $O(Kr)$ (with K being the string tension) between YM monopole and antimonopole.

As to Higgs vacuum BPS monopole solutions, they carry free electric charges in the Higgs phase [22].

Herewith in the concrete case of the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac, the topologically trivial Higgs BPS monopole modes contribute to the "electric" energy (1.11) [5] of the appropriate BPS monopole vacuum, induced by its topological rotations.

There is an interesting and somewhat curious occurs with topologically trivial Higgs vacuum BPS monopoles in the Higgs phase.

On the one hand, they induce vacuum "electric" monopoles (1.13). Herewith the term "electric" is highly relative in this context since this term is associated with temporal vacuum YM components Z_a [16], (1.9), solutions to the Gauss law constraint (1.8), having a definite (rather mathematical) analogy with temporal components, $F_{0\mu}$, of Maxwell tensors, i.e. with electric fields $\mathbf{E}$.

On the other hand, in the Higgs phase [22], topologically trivial Higgs vacuum BPS monopoles $\Phi_{(0)}^a$ carry free electric charges, that are told immediately to vacuum "electric" monopoles (1.13).

Thus vacuum "electric" monopoles (1.13) become indeed *electric* fields in the Higgs phase.

It should be distinguished indeed between the conception of the Higgs phase understood customary (e.g. in Ref. [22]) and that in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac [4].

First of all, the conception of the Higgs phase, as it is understood in [22], is based on fixing *the maximum Abelian gauge* (MAG) and on final violating (in the simple YM case) the initial $SU(2)$ gauge symmetry in the

$$SU(2) \rightarrow U(1) \rightarrow \mathbf{Z} \quad (5.2)$$

wise, involving the "discrete multiplier" $\mathbf{Z}$.

Now let us discuss briefly both these features of the Higgs phase (as it is understood in modern physics).

Fixing MAG comes to singling out the maximum Abelian ($\sim U(1)$) subgroup in $SU(2)$.

The simple way to do this is associated with the diagonal Pauli matrix τ_3 . Then, τ_3 would be considered as the generator of the maximum Abelian subgroup in $SU(2)$. This implies maximizing the integral

$$\int d^4x [(A_\mu^1)^2 + (A_\mu^2)^2] ; \quad A_\mu = A_\mu^a \tau_a, \quad (5.3)$$

or, that is equivalent (see e.g. Ref. [25]), maximizing the quantity

$$\sum_x \sum_{\mu=1}^4 \text{Tr}[\tau_3 U_\mu(x) \tau_3 U_\mu^\dagger(x)]. \quad (5.4)$$

On the other hand, the said is equivalent to decomposing an YM field into its off-diagonal and diagonal components [26]:

$$\mathcal{A}_\mu = A_\mu^a \tau_a + A_\mu \tau_3; \quad a = 1, 2. \quad (5.5)$$

Similarly, one find

$$\mathcal{F}_{\mu\nu} = F_{\mu\nu}^a \tau_a + F_{\mu\nu} \tau_3$$

for the field (YM) strength with off-diagonal and diagonal parts given by

$$F_{\mu\nu}^a = D_\mu^{ab} A_\nu^b - D_\nu^{ab} A_\mu^b$$

and

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu,$$

respectively.

For the YM action (in the Euclidean space E_4) one get finally

$$S_{\text{YM}} = \int d^4x (F_{\mu\nu}^a F_{\mu\nu}^a) + F_{\mu\nu} F^{\mu\nu}.$$

MAG can be got [26] by imposing the Lorentz gauge

$$D_\mu^{ab} A_\mu^b = 0.$$

MAG in YM theories can be generalized easy for the case of the initial $SU(3)_{\text{col}}$ gauge symmetry group in "realistic" QCD models, violated then in the

$$SU(3)_{\text{col}} \rightarrow U(1) \otimes U(1) \quad (5.6)$$

wise.

In this case MAG can be fixed by singling out the Gell-Mann matrices λ_3 and λ_8 from the total set of the Gell-Mann matrices λ .

By analogy with (5.3), now MAG fixing is equivalent to maximizing the integral

$$\int d^4x [(A_\mu^1)^2 + (A_\mu^2)^2 + (A_\mu^4)^2 + (A_\mu^5)^2 + (A_\mu^6)^2 + (A_\mu^7)^2]; \quad A_\mu = A_\mu^a \lambda_a. \quad (5.7)$$

However to achieve the quark confinement in the YM theory and especially in "realistic" QCD models, involving the initial $SU(3)_{\text{col}}$ gauge symmetry group (violated then in the (5.6) wise), the appropriate Abelian $U(1)^{n-1}$ ($n = 2, 3$) subgroups would be also *broken down farther* (and this occurs indeed if MAG is fixed, that results the inevitable violating initial $SU(n)$ gauge symmetries down to their Abelian subgroups with appearance of Higgs modes).

For the initial $SU(2)$ gauge symmetry group this occurs in the (5.2) wise, while for the initial $SU(3)_{\text{col}}$ gauge symmetry group, it is

$$SU(3)_{\text{col}} \rightarrow U(1) \otimes U(1) \rightarrow \mathbf{Z}. \quad (5.8)$$

The said can be grounded by the reasoning to provide the needs of the quark confinement in the above theories.

The sure sign of the quark confinement taking place is, at is well known, the *area law* [24] satisfied by *Wilson loops*, taking the typical shape

$$W(C) \sim \exp\{-KA(C)\}, \quad (5.9)$$

implicating the areas $A(C)$ of surfaces enclosed by appropriate paths C .

Just in order implementing the area law (5.9), characterizing the confinement of quarks and gluons, the phenomenological picture in which quark q ($\bar{q}$) and gluons A are confined by (infinitely thin) Meissner flux tubes is quite fit.

This phenomenological picture is similar to that proper to superconductors.

In particular, in superconductors, the magnetic field $\mathbf{B}$ decays exponentially at penetrating into the superconducting region [31]:

$$\mathbf{B}(x) = \mathbf{B}_0 e^{-x/\lambda_L}, \quad (5.10)$$

with

$$\lambda_L = \frac{1}{M_\gamma} \quad (5.11)$$

being the *penetration or London depth*, depending on the "photon mass"

$$M_\gamma = \sqrt{2} ea. \quad (5.12)$$

The parameter a appears in latter Eq.

It plays the role of the order parameter in the *Ginzburg-Landau* superconductivity model [30, 31, 36]: $a = 0$ for the normal phase, while $a \neq 0$ for the superconducting phase.

The magnetic field $\mathbf{B}$, (5.10), decaying exponentially at penetrating into the superconducting region, is the solution to the *Maxwell-London equation* [31, 36]

$$\text{rot } \mathbf{B} = \text{rot rot } \mathbf{A} = \mathbf{j}_s = 2e^2 a^2 \mathbf{A}, \quad (5.13)$$

implicating the preserved Nöther current [13, 36]

$$\mathbf{j}_s = -i \frac{e}{m} (\phi^* \nabla \phi - \phi \nabla \phi^*) - \frac{e^2}{m} |\phi|^2 \mathbf{A}, \quad (5.14)$$

with ϕ being the superconductor wave function.

Eq. (5.10) serves as the mathematical expression for the Meissner effect. The essence of this effect is in expulsion of the magnetic field from the superconducting region.

Similar locating the "magnetic" flux in the shape of infinitely narrow tubes explains the confinement effect in QCD.

The *confinement phase* can be treated [22] as dual to the Higgs phase, discussed above.

In this phase gluons and quarks, that are purely electric objects, are confined by infinitely narrow "magnetic" flux tubes (strings). Herewith Higgs fields become free and purely magnetic objects.

The occurred in the confinement phase requires undoubtedly the spontaneous breakdown of the initial ($SU(3)_{\text{col}}$ or $SU(2)$) gauge symmetries groups with the appearance of Higgs modes.

In this case also MAG fixing seems to be quite natural.

In the modern literature such (phenomenological) picture of confinement based on MAG fixing and the dual superconducting picture (similar to that one can observe in the Ginzburg-Landau model [30] and implicating [22] the Higgs and confinement phases) is referred to as the *Abelian dominance* [32]).

But, indeed, it is not sufficient to violate the initial $SU(n)$ ($n = 2, 3$) gauge symmetry group in a non-Abelian model up to its $U(1)^{n-1}$ subgroup in order to ensure the quark confinement in that model.

The thing is that the appearance of (infinitely thin) "magnetic" flux tubes (strings) confining quarks in QCD can be explained good in the framework of the well-known NO model [27].

This model serves as a highly correct representation for the Abelian Higgs theory (to which the non-Abelian Higgs theory reduces upon MAG fixing).

With the example of the initial $U(1)$ gauge symmetry group afterwards violated, in the original paper [27] it was demonstrated the existence of specific solutions to the equations of motions, *Nielsen-Olesen vortices* (see e.g. [31, 36]).

The way grounding the existence of NO vortices in the Abelian Higgs theory comes to violating the initial $U(1)$ gauge symmetry group proper to that theory (this involves the nonzero value $|\phi| \neq 0$ for the Higgs field ϕ in the "asymmetric" phase, entering explicitly [31, 36] the appropriate equations of motions) and, from the topological standpoint, to the presence of thread topological defects (just interpreted as NO vortices) inside the appropriate vacuum manifold.

In the light of the said, the task to ascertain the look of this vacuum manifold becomes very important.

The investigations about the NO model [27] are not the immediate goal of the present study, but some outlines and conjectures concerning this model will be helpful for us in understanding of processes taking place in QCD if stand on the Abelian dominance viewpoint [32].

From our previous discussion about the liquid helium II theory, some analogies suggest concerning the origin of NO vortices in the Abelian Higgs model [27].

First of all, from the topological point of view, NO vortices, as a particular case of thread topological defects, would be associated with the "discrete" vacuum geometry should be assumed for the appropriate degeneration space.

Such degeneration space (vacuum manifold) can be obtained in the way similar to that in which the discrete vacuum manifold $\tilde{R}$ has been got in the rested liquid helium II specimen case.

And again, as in the rested liquid helium II specimen case, the degeneration space in the NO model [27], coincides to within an isomorphism with the quite separate set we have denoted above as $\tilde{U}$ (on the other hand, the alone this discrete set $\tilde{U}$ can be treated as the residual gauge symmetry group in the NO model [27]).

However there are some principal distinctions between the latter case and the Abelian Higgs model [27] involving NO vortices.

So, for instance, the first-order phase transition occurs primary in that model (see the arguments [28], repeated also in Ref. [6]). It comes to a discontinuity in the plot of the appropriate order parameter (as a such order parameter, the vacuum expectation value $\langle |\phi|^2 \rangle$ for the Higgs field ϕ can be chosen). This implies the coexistence of the metastable (symmetric) and stable (asymmetric) thermodynamic phases in the Abelian Higgs model [27].

This discontinuity in the plot of $\langle |\phi|^2 \rangle$ implies (cf. [29]) the finite difference $\Delta F(\langle |\phi|^2 \rangle)$ of the appropriate free energy, interpreted as the latent heat.

It is just the supercooling case (according to the argument similar to those [41] we have encountered above at the analysis of vortices in a liquid helium II specimen).

Once thus, the way (3.24) violating the $U(1)$ gauge symmetry (fit for the *second-order phase transition* [46] in a helium specimen) is not quite suitable in the Abelian Higgs model [27] with NO vortices implying the *first-order phase transition*.

In the first case, the symmetric thermodynamic phase disappears entirely upon $U(1)$ violating.

This makes quite correct Eq. (3.24) for the quite separate set [40] $\tilde{U}$, coinciding with the degeneration space (vacuum manifold) $\tilde{R}$.

The look of this discrete space (in which all the topological domains refer to the asymmetric thermodynamic phase) just represents the above (primary) second-order phase transition in the helium theory.

Unlike the former case, in the Abelian Higgs model [27], in which the first-order phase transition occurs involving coexisting [6, 28] the stable and metastable thermodynamic phases, it can be assumed $U(1)$ violating in such a wise that a part of topological domains in the $U(1)$ group space remains "glued together", representing herewith the "symmetric" (metastable) thermodynamic phase, while the rest of topological domains become a discrete (quite separate) set, representing the asymmetric (stable) thermodynamic phase.

Indeed, this "coexistence" of the metastable and stable thermodynamic phases in the Abelian Higgs model [27] with NO vortices cannot last infinitely long time.

After all (at lowering the temperature below the Curie point T_c : to some its value $T_b < T_c$, or under influence of external or internal fluctuations), the Higgs field ϕ in the Abelian model [27] leaving the "false" vacuum $|\phi| = 0$ downs rapidly into its "true" minimum (say, $|\phi| = \phi_0 \neq 0$), and the metastable (symmetric) thermodynamic phase (corresponding to the above described continuous set and the "false" vacuum $|\phi| = 0$) disappears herewith entirely.

From the "geometrical" standpoint, the gradual destruction of the continuous $U(1) \simeq S^1$ group space (with forming the "bubble" [28] of the new, stable and "discrete", thermodynamic phase inside the metastable, "continuous", one) occurs.

At this disappearing the metastable (symmetric) thermodynamic phase, the accumulated latent heat is released; it is just the *reheating* situation [29] ²⁰.

It may be guessed that "final" downing the Higgs field ϕ in the "true" minimum ϕ_0 corresponds to the dual superconducting picture of confinement in QCD based upon MAG fixing and the Abelian Higgs model [27] with NO vortices.

In connection with our discussion about the first-order phase transition taking place in the Abelian Higgs model [27] (implicating NO vortices), it will be relevant to mention the very important analogy (in the sphere of the topology and phenomenology) between this model and the case [13, 30, 31] of Type II superconductors.

In Type II superconductors, the first-order phase transition occurs with definite conditions in the presence of an background magnetic field $\mathbf{H}$.

First of all, it would be [13, 30, 31] $\kappa > 1/\sqrt{2}$ for the Ginzburg-Landau parameter

$$\kappa = \frac{\lambda_L}{\xi},$$

with λ_L given in (5.11) [31] and ξ being the coherence length whose physical sense is [13] the correlations radius of fluctuations in the order parameter $< |\phi|^2 >$.

Generally speaking [13], the Ginzburg-Landau parameter κ is a function of the temperature T : $\kappa \equiv \kappa(T)$.

As it was demonstrated in the original paper [30] (see also [13]), the inequality $\kappa < 1/\sqrt{2}$, true for Type II superconductors, is associated with the negative surface tension $\alpha_{ns} < 0$ (which look is (4.22)) between the normal and superconductor phases.

In the latter case it becomes advantageous, from the thermodynamic standpoint, to compensate increasing the volume energy by this negative surface energy $\alpha_{ns} < 0$.

This implies arising germs of the n-phase inside the s one at the values H of the background magnetic field $H = |\mathbf{H}|$ exceeding a value H_{c1} , called [13] *the lower critical field*.

Vice verse, germs of the s-phase inside the n one arise at the values H don't exceeding a field H_{c2} , called *the upper critical field*.

Thus in the interval $H_{c1} < H < H_{c2}$, one can observe [13] the *mixed state* in Type II superconductors (there are definite alloyed metals and combinations of metals). In this case a Type II superconductor abides simultaneously in the n and s states.

At $H \leq H_{c1}$, a specimen is purely in the s-state, while it is purely in the n-state at $H \geq H_{c2}$.

Indeed, as it was argued in [53] (see ibid §56), the value of the lower critical field H_{c1} is specified by the geometry of the tested superconductor specimen. Mathematically, this

²⁰This latent heat is equal quantitatively to the magnetic energy

$$\sim \int d^3x B^2,$$

associated with NO vortex solutions.

implies that $H_{c_1} = (1 - n)H_{c_2}$, with n being now a *demagnetizing coefficient* ²¹.

It is easy to understand that the mixed state in Type II superconductors is the sign of the first-order phase transition occurring therein.

Herewith it is again the typical supercooling case when the symmetric n-phase is the metastable and asymmetric s-phase is the stable phase (cf. the above discussed supercooling case in the NO model [27] in the temperature interval $[T_b, T_c]$; there is the only correction that now the situation is developed towards the *restoration* of the $U(1)$ gauge symmetry proper to the quested model) ²².

It is enough manifest that the case of Type II superconductors can be treated as a nonrelativistic limit of the Abelian Higgs NO model [27] ²³.

As in the NO model [27], the nontrivial topological content of the Ginzburg-Landau superconductivity theory [30] can be discovered.

This can be seen at writting out explicitly the gauge ($U(1)$) invariant superconductivity current [13]

$$\mathbf{j}_s = \frac{\hbar e}{2m} n_s (\nabla \Phi - \frac{2e}{\hbar c} \mathbf{A}), \quad (5.15)$$

²¹The computations [53] show that $n = 1/3$ for a ball; thus in this case $2/3 H_{c_2} < H < H_{c_2}$. For a cylindrical specimen in a transverse magnetic field, $n = 0.5$ and $0.5 H_{c_2} < H < H_{c_2}$. In the longitudinal magnetic field, $n = 0$, thus $H_{c_1} = H_{c_2}$, and the mixed state is absend. This means that superconductivity is destroyed completely immediately at a fixed H_c . Finally, for a plane-parallel plate in a transverse magnetic field $n = 1$, and it is in the mixed state at any $H < H_{c_2}$.

²²The correlation between the interval $H_{c_1} < H < H_{c_2}$ for the background magnetic field H in the mixed state and the temperature T is, indeed, enough complicate (see e.g. [13]). In spite of this, it is obvious that in the Curie point T_c , $H_{c_1} \rightarrow H_{c_2} \rightarrow 0$. The latter fact can be explained [53] by the thermodynamic reasoning. Really, the experimental data tesify that a second-order phase transition occurs in a superconductor in the absence of background magnetic fields.

On the other hand, it is well known from the general theory of second-order phase transitions that a change of the thermodynamic potential $\Phi(p, T)$ near the aproprate Curie point T_c is directly proportional to $(T - T_c)$.

Then we take into account [13, 53] that

$$\Phi_s(p, T) + \frac{H_c^2}{8\pi} V_s(p, T) = \Phi_n(p, T),$$

where $V_s = \partial \Phi_s(p, T) / \partial p$.

Since the plot of $\Phi(p, T)$ is comtinuous in the Curie point T_c at a second-order phase transition if occurs in a superconductor, it should be $\Phi_s(p, T) = \Phi_n(p, T)$ in that case. Whence one can conclude [53] that

$$H_c = \text{const}(T - T_c)$$

in an infinite neighbourhood of T_c .

The said allows to construct the plots $H(T)$ for the s, n and mixed states in Type II superconductors (see Fig. 7 in [13]).

Note by the way that for a cylinder located in a longitudial background magnetic field (when $n = 0$), also a second-order phase transition occurs.

²³It is so at identifying [31, 36] the density of a superconducting *Cooper pair* with the Higgs field squared $|\phi|^2$ in the NO model [27].

involving the value n_s , the density of superconductive electrons.

Introducing such a current corresponds to the gauge transformation law

$$\Phi \rightarrow \Phi + \frac{2e}{\hbar c} \chi(\mathbf{r}) \quad (5.16)$$

for the phase Φ of the superconductor wave function ϕ (with the charge $2e$ coming from the presence of Cooper pairs in the Ginzburg-Landau superconductivity theory [30]).

Simultaneously, electromagnetic potentials $\mathbf{A}$ would be suffered standard $U(1)$ gauge transformations

$$\mathbf{A} \rightarrow \mathbf{A} + \nabla \chi(\mathbf{r}) \quad (5.17)$$

in order for the Hamiltonian in the Ginzburg-Landau theory [30] to be manifestly gauge invariant.

Note that Eq. (5.15) or the superconductivity current is the alternative look of Eq. (5.14) [36]. The way going over from (5.15) to (5.14) comes to applying the normalization [13]

$$\phi = \sqrt{\frac{n_s}{2}} e^{i\Phi}.$$

In the light of Eq. (5.15), nontrivial topologies in the Ginzburg-Landau superconductivity theory [30] originate from the Maxwell-London equation (5.13) [13, 36], implicating the nonzero curl

$$\text{rot } \mathbf{j}_s = -\frac{e^2 n_s}{mc} \mathbf{B} \quad (5.18)$$

(indeed, the latter result can be got immediately from (5.15)).

Comparing Eq. (5.18) with the results (for instance, (4.9)) in the liquid helium II at rest theory [15], one concludes about the presence of topologically nontrivial vortices in the Ginzburg-Landau superconductivity theory [30].

More exactly, since $\text{rot } \mathbf{j}_s \neq 0$ comes to $\mathbf{B} = \text{rot } \mathbf{A} \neq 0$, it is just the source of nontrivial topologies $n \neq 0$ in the Ginzburg-Landau theory [30]. Herewith the specific of the mentioned superconductivity theory is that such nontrivial topologies are compatible with the superconductivity effects in that theory.

This distinguishes principally the Ginzburg-Landau superconductivity theory [30] from the liquid helium II (at rest) theory, in which the superfluidity is associated namely with the trivial topology $n = 0$. The said is the purely gauge effect connected with the presence of electromagnetic potentials $\mathbf{A}$ in the superconductivity theory [30].

Choosing an arbitrary closed integration way C , we can write down for $\mathbf{A}$:

$$\oint_C \mathbf{A} d\mathbf{l} = \int \text{rot } \mathbf{A} \cdot d\mathbf{f} = \int \mathbf{B} d\mathbf{f} \equiv \Phi,$$

where in this context, Φ is the flux of the magnetic field $\mathbf{B}$.

In particular, when the integration way C is chosen to lie inside a superconductor far from its surface (where the superconductivity current $\mathbf{j}_s$, (5.15), runs actually), $|\mathbf{j}_s| \rightarrow 0$ due to (5.10), (5.18).

This implies

$$\oint_C \mathbf{A} d\mathbf{l} = \frac{\hbar c}{2e} \oint_C \nabla \Phi \cdot d\mathbf{l} = \frac{\hbar c}{2e} \delta \Phi. \quad (5.19)$$

where $\delta \Phi$ is the change of the phase Φ at the complete round along the integration way C .

However from the claim for the superconductor wave function to be finite, it follows that this change of the phase Φ can take only values $2\pi n$ ($n \in \mathbf{Z}$).

One comes thus to the result [13]

$$\Phi = n\Phi_0, \quad \Phi_0 = \frac{\pi \hbar c}{|e|} = 2 \cdot 10^{-7} \text{ G cm}^2 \quad (5.20)$$

for the phase Φ . The value Φ_0 is called *the elementary quant of the magnetic flux* [13].

In Eq. (5.19) the variation $\delta \Phi$ is indeed infinitesimal, that corresponds to the fact that $\text{rot } \nabla \Phi = 0$. This can be checked also immediately recalling [13] that all the values in the Ginzburg-Landau superconductivity theory [30] are assumed to be distributed to depend only on x , the depth of the superconductor, while four-potentials A are taken in the *London* gauge $\text{div } \mathbf{A} = 0$; $A_0 = 0$; $A_x = 0$.

In this case ²⁴

$$|\mathbf{B}| = B_z = A' \equiv dA/dx,$$

i.e.

$$A = A_y \sim \int |\mathbf{B}| dx \sim B_0 \int e^{-x/\delta} dx = -\delta B_0 e^{-x/\delta} \rightarrow 0, \quad x \rightarrow \infty.$$

that proves the above statement (at choosing the way C to lie in the (x, y) -plane far from the superconductor surface).

Thus one can observe actual disappearing nontrivial topologies deep into a superconductor.

The patterns of Type II superconductors are superconducting alloys and polymetallic connections [13].

Say also several words about Type I superconductors, for which $\kappa < 1/\sqrt{2}$ and, respectively, $\alpha_{ns} > 0$. The patterns of such superconductors are pure metals [13].

In Type I superconductors first-order phase transition occurs when $H < H_c$. Meanwhile, the role of the surface tension in Type I superconductors comes to hampering somehow forming germs of the s-phase.

And moreover, the s-phase is maintained as a metastable one at magnetic fields $H > H_c$. Simultaneously, at $H < H_c$ the n-phase disappears completely; thus [13] there is no mixed state in Type I superconductors ²⁵.

²⁴According to the geometrical reasoning, $\mathbf{A}$ would then lie in the (y, x) -plane because locating the magnetic tension $\mathbf{B}$ in the (y, z) -plane perpendicular to the axis x , the direction deep into the superconductor.

This implies $A_y \equiv A$ and therefore $|\mathbf{B}| = B_z = A' \equiv dA/dx$.

²⁵However, as it was demonstrated in [13], if κ varies in the interval

$$1/\sqrt{2} = 0.71 > \kappa > 0.59/\sqrt{2} = 0.42,$$

Indeed, Eq. (5.19), implicating $|\mathbf{j}_s| \rightarrow 0$ as $|x| \rightarrow \infty$, is correct [13] only in the absence of the external magnetic field H , that corresponds [53] to the second-order phase transition. If such a magnetic field is switched on, i.e. when the first-order phase transition occurs actually in a superconductor, the superconductivity in a mixed state is violated (in a cylindrical specimen) along the *Abrikosov threads* [49], directed, in turn, along the symmetry axis (z) of this cylindrical specimen. The cores of such Abrikosov threads have the effective radii $\approx \xi$.

As a result [13], superconductivity currents form rings in the (y, z) -plane perpendicular to the superconductor depth x , while Abrikosov threads serve as “embryos” of the n-phase.

If we denote now as $r(x)$ the distance from an Abrikosov thread ($r = 0$ for this thread), then [13]

$$j(n)_s = \frac{c}{4\pi} |\text{rot} \mathbf{B}| = \frac{c\Phi_0}{8\pi^2\delta^2 r} n, \quad n \in \mathbf{Z}. \quad (5.21)$$

Simultaneously [13], there is the following dependence between the topological number n which an Abrikosov thread possesses and the magnetic induction $\mathbf{B}$ which this thread carries:

$$B(r) = \frac{\Phi_0 n}{2\pi\delta^2} \ln \frac{\delta}{r}; \quad \xi \leq r \leq \delta. \quad (5.22)$$

Actually, this value $B(r)$ of the magnetic induction refers to an (local) thermodynamin equilibrium state in the given Type II superconductor. This thermodynamin equilibrium is set course the first-order phase transition occurring therein.

This reflects the general nature of thread topological defects in models implicating first-order phase transitions. Although [9] the (local) thermodynamin equilibrium is violated in the coordinate regions where thread topological defects are located, in models implicating first-order phase transitions such an equilibrium is then restored again. This involves, in turn, items in the appropriate Hamiltonians describing thread topological defects.

In particular, for Type II superconductors it is [13] the $\sim L\epsilon$ item (with L being the length of the thread and ϵ being the energy density per the length unit of this thread). For the liquid helium II rigidly turning model, it is the item given by Eq. (3.18) [13].

Thus we see that the superconductivity currents $j(n)_s$ are topologically degenerated in the presence of Abrikosov threads carrying the magnetic field $\mathbf{B}(r)$ (related as $\mathbf{B} = \mathbf{H} + 4\pi\mathbf{M}$ to the external magnetic field $\mathbf{H}$). It is quite plausible herewith locating such currents inside the superconductor (i.e. at arbitrary x). It can be seen from (5.21).

mixed states can appear also in Type I superconductors.

Herewith at $H_{c_2} = \sqrt{2} \kappa H_c$ one can find such value

$$H_{c_3} = 1.7H_{c_2} = 2.4\kappa H_c$$

of the magnetic tension H at which *surface* germs of the s-phase arise.

And moreover, in the interval $H \in [H_{c_2}, H]$, the n-phase becomes metastable, while it is completely unstable as $H < H_{c_2}$. Thus the mixed state arises in Type I superconductors at the above specific conditions [13] imposed onto the Ginzburg-Landau parameter κ and the external magnetic field H in connection with the mentioned above fact maintaining the s-phase as metastable in Type I superconductors at some values $H > H_c$.

In this (as it was stressed, for instance, in the review [31]) is just the similarity between NO vortices [27] and Abrikosov threads in Type II superconductors [30]. Both are patterns of thread topological defects corresponding to the spontaneous $U(1)$ breakdown course the first-order phase transition occurring.

In the NO model [27] the role of the Ginzburg-Landau parameter κ is played [31] by

$$\kappa = \sqrt{\beta}, \quad \beta = \frac{\lambda}{2e^2},$$

with λ being the Higgs selfinteraction.

In this case the magnetic flux

$$\Phi_B^n = \int_{\Sigma} B d^2x = \oint_{\mathcal{C}} \mathbf{A} \cdot d\mathbf{s} = n \frac{2\pi}{e}, \quad (5.23)$$

taking over the surface Σ bounded by an asymptotic curve $\mathcal{C}$, where the magnetic field $\mathbf{B}$ is generated by NO vortices solutions [27, 31, 36] (in the Higgs and gauge sectors of the NO model), is similar, when $\sqrt{\beta} > 1/\sqrt{2}$, to the magnetic flux $n\Phi_0$, (5.21), crossing [13] the (cylindrical) specimen, in the case of Type II superconductors.

There is, however, the principal distinction between vortices [13, 15] in a liquid helium II (turning) specimen and those (Abrikosov threads) arising in a Type II superconductor [13, 49].

In the liquid helium II (turning) case topologically nontrivial vortices belong to the appropriate vacuum manifold $\tilde{U}(1)$, topologically equivalent to the discrete set $\mathbf{Z}$ of integers. In the Type II superconductors case there are two types of (topologically nontrivial) vortices. Some of them (just referring to as Abrikosov threads [49]) belong to the n component in the investigated specimen, i.e. to the symmetrical thermodynamic phase. It is so due to our above hypothesis that a first-order phase transition occurs in Type II superconductors at which, at $T \rightarrow 0$, the “symmetrical” thermodynamic phase, representing by remaining “gluing together” topological domains coexists with the “less symmetrical” thermodynamic phase, representing by a subset of the discrete set $\mathbf{Z}$. The second type of vortices are those existing inside the appropriate discrete vacuum manifold, that is again $\tilde{U}(1) \simeq \mathbf{Z}$ ²⁶.

This “status quo” between the mentioned thermodynamic phases “freezes” (see below) at such low temperatures ²⁷.

²⁶The same assertion about two kinds of vortices is correctly also for the NO model [27], possessing the similar geometrical features and in which the first-order phase transition proceeds in the same wise.

²⁷The fact that Abrikosov threads belong to the “symmetrical” n component of a Type II superconductor allows (see the original paper [49]) to set $|\phi(0)|^2 = 0$ for the superconductivity wave function $\phi(r)$ squared (actually, it should be set within the $|r| \leq \xi$ neighborhood of $r = 0$, i.e. within a typical Abrikosov thread core), serving naturally as an order parameter in the superconductivity model. The role of $\phi(r)$ as an order parameter in the superconductivity model is similar to the role of the order parameter $|\Xi(t, \mathbf{r})|^2$, given via Eq. (1.4) [13], in the liquid helium II model. However, there is a distinction between the both order parameters.

In the second case, as it was shown above repeating the arguments [41], the value $|\Xi(t, \mathbf{r})|^2$ diverges

Eq. (5.21) [13] allows to refine the topological structure of the both thermodynamic phases in Type II superconductors and to suggest a definite hypothesis about the way violating the $U(1)$ gauge symmetry therein.

We see from (5.21) that an Abrikosov vortex possessing the topological number n and belonging to a “continuous and symmetrical” n-layer, induces the superconductivity current $j(n)_s$, belonging to a “discrete and less symmetrical” s-layer. This indicates the close correlation between the above described continuous and discrete sets. And moreover, it can be suggested that the discretization process inside the $U(1) \simeq S^1$ group space in Type II superconductors proceeds in such way that within a topological domain possessing a topological number n , a domain wall (surmounted for the magnetic field $\mathbf{B}$) arises between topologically equivalent loops. Thus, geometrically and thermodynamically, the S^1 circle coexists in Type II superconductors with a discrete set, topologically equivalent to $\mathbf{Z}$.

Note that arising circular superconductivity currents $j(n)_s$, (5.21), in Type II superconductors induced by Abrikosov threads, belonging to n-layers therein, is, probably, in a good agreement with the London’s postulate (1935 y.) about the existence of a finite energy slit between n and s states.

The above suggestion about the discretization process inside the $U(1)$ group space can prove to be plausible also in the NO model [27], just as the opinion that such process proceeds toward splitting whole topological domains from the continuous $U(1) \simeq S^1$ space.

The outlined above parallel between the Type II superconductivity and the Abelian Higgs NO model [27] promotes also the comprehension of the dual superconducting picture [22, 31] of the confinement of quarks and monopoles.

One can imagine [31], in his thoughts, placing some ”north” and ”south” magnetic monopole inside a Type II superconductor in such a wise that they are separated by the distance d .

Thinking that the (background) magnetic field $\mathbf{B}$ is concentrated in cores of vortices

logarithmically in points where vortices are located, and this can be treated as an evidence in favour of the first-order phase transition occurring in a liquid helium II specimen.

In the first case (see e.g. §48 in [13]) the plot of $|\phi(r)|^2$ falls rapidly to zero in the $|r| \leq \xi$ neighborhood of $r = 0$, remaining almost constant at other values of r , corresponding to the s state. One can speak in this case that such a plot is not smooth at $r = 0$. It proves to be once again the argument in favour of the first-order phase transition occurring in Type II superconductors. Roughly, the reasoning comes to the following. The free energy F of a superconductor contain always the item [13]

$$\int dV \frac{\hbar^2}{4m} |\nabla \phi|^2$$

Proving precisely that this integral diverges because $|\nabla \phi|^2$ does not exist in points corresponding to locating Abrikosov threads, one can conclude that the plot of the free energy F is not continuous and whence also the entropy $S = (E - F)/T$ of a Type II superconductor can suffer a discontinuity $\Delta S = S_n - S_s$ if F diverges faster than T in the $T \rightarrow 0$ limit. It is just the sign of the first-order phase transition occurring.

and will not extend into the superconducting region, the field energy of this system becomes [31]

$$V = \frac{1}{2} \int d^3x \mathbf{B}^2 \propto \frac{4\pi d}{e^2 \lambda_L^2}. \quad (5.24)$$

Thus, the interaction energy of magnetic monopoles (quarks) grows linearly with their separation d , as it would be in the dual superconducting picture [22] of confinement.

The Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac [4] results the approach to QCD (including the confinement picture) rather another that one gets fixing MAG and assuming the dual superconducting picture [22] of confinement (implicating the Mandelstam linearly increasing potential).

Firstly, as it was discussed in Refs. [5, 12] (see also [1]), at constructing Minkowskian Gauss-shell QCD (generalizing the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac and involving the initial $SU(3)_{\text{col}}$ gauge symmetry group), the specific gauge for the "intermediate" $SU(2)_{\text{col}}$ symmetry group in the "breakdown chain"

$$SU(3)_{\text{col}} \rightarrow SU(2)_{\text{col}} \rightarrow U(1)$$

can be fixed.

It comes [1, 5, 12] to the choice of the antisymmetric Gell-Mann matrices $\lambda_2, \lambda_5, \lambda_7$ as the generators for $SU(2)_{\text{col}}$.

At further assuming the discrete geometry (3.3) for $SU(2)_{\text{col}}$, this involves lot of consequences for Gauss-shell QCD (described briefly in Ref. [1]).

Among such consequences fixing the $\lambda_2, \lambda_5, \lambda_7$ ("antisymmetric") "gauge", one can mention fermionic rotary (axial) degrees of freedom $\mathbf{v}_1 = \mathbf{r} \times \mathbf{K}$ (with $\mathbf{K}$ being the polar colour vector, $SU(2)_{\text{col}}$ triplet).

Repeating the arguments [17], the source of the above fermionic (quark) rotary degrees of freedom can be found in the interaction item $\sim Z^a j_{Ia(0)}$ in the Gauss-shell reduced Hamiltonian of constraint-shell QCD.

This item implicates the zero-mode solution Z^a , (1.9), to the Gauss law constraint (1.6) and the fermionic charge

$$j^{Ia(0)} \sim g \bar{\psi}^I (\lambda^a/2) \gamma^0 \psi^I.$$

Here it would be set $a = 2, 5, 7$ and γ^0 is the Dirac matrix; $\bar{\psi}^I$ and ψ^I [20] are fermionic Dirac variables involving "small" Gribov multipliers $v^{(0)}(\mathbf{x})$.

Thus the existence of fermionic (quark) rotary degrees of freedom $\mathbf{v}_1$ (similar somewhat to rotary terms in molecules) can be associated lawfully with the "discrete vacuum geometry" assumed for Minkowskian constraint-shell QCD in the Dirac fundamental scheme [4].

Secondly, the Mandelstam linearly increasing potential $\sim Kr$ [24] in Minkowskian constraint-shell QCD "draws back in a background".

More exactly, the Dirac fundamental scheme [4] for Minkowskian constraint-shell QCD implicates a linear combination of well-known instantaneous ("four-fermionic") interaction

potentials: Coulomb and "golden section" ones [1, 5, 12, 20] (with coefficients depending somehow on the temperature T and flavours mass scale m) as the source of hadronization and confinement in that model (see e.g. the papers [12, 54, 55]).

Such linear combination may be got at finding the Green function of the Gauss law constraint equation in the BPS (Wu-Yang) monopole background ²⁸.

The important objection restricted the role of the Mandelstam linearly increasing potential $\sim Kr$ in QCD was given also in Ref. [57].

It turns out that for heavy quarkonia, only the Coulomb potential $\sim 1/r$ contributes in QCD.

Then the problem arises, to link (continuously and smoothly) the Mandelstam linearly increasing and Coulomb potentials in order to satisfy the heavy quarkonia limit [57].

As to the Mandelstam linearly increasing potential $\sim Kr$ [24], the hypothesis can be suggested about including this potential at (finite) temperatures $T \neq 0$ in the linear combination with the Coulomb and "golden section" potentials. This can be done via a selection of the proper coefficient for the Mandelstam potential.

Among consequences for Minkowskian constraint-shell QCD of such including one can point out the "ordinary" annihilation mechanism [48] for monopole-antimonopole pairs via Abrikosov threads [9, 49] of the "magnetic" field (this mechanism was us discussed above).

This is correlated closely with the dual superconducting picture of confinement.

More exactly, at placing a monopole-antimonopole pair into the hot YM plasma, the typical length of the Abrikosov thread ("magnetic" tube) between this pair becomes [28, 48] $\Delta l \sim (g^2 T)^{-1}$. This implicates the temperature-depended Mandelstam potential in the shape $K(g^2 T)^{-1}$.

On the other hand, this including the temperature-depended Mandelstam potential in Minkowskian constraint-shell QCD can give rise to series of problems and questions.

In particular, the problem is to check satisfying the area law (as the confinement criterion) in the situation of specific "**Z**-dominance" taking place actually in Minkowskian constraint-shell QCD and induced by **Z**-vortices (thread topological defects) inside the appropriate vacuum manifold, assumed to be discrete and given through (3.9).

Speaking about the "**Z**-dominance" in Minkowskian constraint-shell QCD based upon the Dirac fundamental quantization scheme [4] it should be borne in mind the deep distinction between this "**Z**-dominance" and that one can observe actually at enlisting the Abelian Higgs NO model [27] (involving NO vortices) in order to explain the dual superconducting picture of (quark) confinement in QCD.

In spite of a definite resemblance between the both approaches to QCD, involving thread topological defects (**Z**-vortices) inside appropriate vacuum manifolds, there is indeed a "watershed" between these approaches.

²⁸ Wu-Yang monopoles [56], as solutions to the equations of motions in the "purely YM" model, can be considered as spatial asymptotes for YM BPS monopoles.

This "watershed" is actually the "watershed" between the Dirac fundamental quantization method [4] and the FP "heuristic" one [35]²⁹ in gauge physics.

Discussion.

Completing this study, we should like talk over some problems leaving outboard in the main part of the work.

Firstly, we should like discuss some aspects of first-order phase transitions.

From our previous discussion, the nature of the first-order phase transition occurring in the Abelian Higgs NO model [27] with NO vortices seems to be enough transparent.

At assuming the "discrete vacuum geometry" of the $\tilde{U}$ type, with $\tilde{U}$ being a quite separate set [40], the nature of the first-order phase transition in the Abelian Higgs NO model [27] keeps within the framework of the standard scheme [29] of such phase transitions.

In the liquid helium II (at rest) model [15, 41] the situation is indeed by far complicate.

In this model the second-order phase transition (confirmed [41, 46] by experimental data) occurs primary. As a result, the "asymmetric" liquid helium II phase, possessing manifest superfluid properties, arises (while the "symmetric" liquid helium I phase [46] disappears entirely).

Then the first-order phase transition [41] occurs in a liquid helium II (at rest) specimen and (rectilinear) vortices [15] appear in this specimen.

Here the question about the dynamics of such "double" phase transition arises. And moreover, it is important to co-ordinate this dynamics with the way (3.24) violating the initial $U(1)$ gauge symmetry of the helium Bogolubov Hamiltonian $\hat{\mathcal{H}}$ [41], implicating the "discrete vacuum geometry" $\tilde{U}$ (assuming such "discrete vacuum geometry" explains correctly vortices [15] in a liquid helium II specimen).

The following version can be proposed to attempt answer this question.

The second-order phase transition in the helium (at rest) model occurs with the separation (by a wall) of the topologically trivial domain from the whole $U(1)$ group space.

This topologically trivial domain inside the $U(1)$ group space just corresponds to superfluid potential motions in a liquid helium II specimen.

Then the further destruction of the $U(1)$ group space occurs in the course of the first-order phase transition therein.

²⁹As an (indirect) evidence in favour "belonging" the Abelian ($\mathbf{Z}$) dominance in the spirit [32] to the FP "heuristic" quantization scheme [35] one can point out [22] getting Polyakov hedgehogs

$$a_k(x) \sim (x - x_0)_k,$$

with $a_k(x)$ being the "off-diagonal" gauge field in (5.5), arising at the $SU(2) \rightarrow U(1)$ breakdown when MAG is fixed in the Higgs theory.

On the other hand, Polyakov hedgehogs [34], as a particular case of 't Hooft-Polyakov monopoles [33, 34, 36], are described correctly in the framework of the FP "heuristic" quantization scheme [35] (including [1, 6] fixing the Weyl gauge $A_0 = 0$; at MAG fixing, also the additional $\delta(D_\mu^{ab} A_\mu^b)$ multiplier in the appropriate FP path integral would arise reflecting this Lorenz gauge being assumed [26]).

As a result, the initial $U(1)$ gauge symmetry of the helium Bogolubov Hamiltonian $\hat{\mathcal{H}}$ becomes violated finally in the (3.24) wise with appearing quantum vortices [15].

The next important remark regarding the liquid helium II (at rest) case is "freezing" the supercooling situation at temperatures near the absolute zero $T = 0$.

In this case, honestly speaking, it is not so important, which of thermodynamic phases (the phase of superfluid potential motions or the phase of quantum vortices) would be treated as the stable and which as the metastable.

Analogous "freezing" takes place also in a Type II superconductor specimen which is in the mixed state. In this case the geometrical structure of a Type II superconductor, at which [53] its geometrical structure with an alternation of n, "continuous", and s, "discrete", layers can be maintained infinitely long at maintaining the low temperature $T \rightarrow 0$.

This can be demonstrated analytically repeating the arguments [53]. Ibid it was shown that the difference of entropies $\Delta S = S_n - S_s$ between the (n) and (s) states is indeed

$$\Delta S = -V_s \frac{\partial}{\partial T} \left(\frac{H_c^2}{8\pi} \right),$$

with V_s being the volume of the (cylindrical) specimen.

Then the released (latent) heat of the first-order phase transition taking place in a Type II superconductor at a fixed temperature T will be

$$Q = T \Delta S = -\frac{V_s H_c T}{4\pi} \left(\frac{\partial H_c}{\partial T} \right)$$

at a constant pressure p . It approaches zero as $T \rightarrow 0$ as well as at $H_c \rightarrow 0$. As it was argued above [53] the latter fact corresponds to the Curie point $T = T_c$ since $H_c = \text{const}(T - T_c)$. Thus at $T \rightarrow 0$ the both, (n) and (s), thermodynamic phases freeze, coexisting each with other.

In the connection with the said, there is worth to recall that the above relations were derived in the monograph [53] issuing from the equality $\mathcal{F}_s = \mathcal{F}_n$ for the specific thermodynamic potentials $\mathcal{F}_s$ and $\mathcal{F}_n$ (per an unit volume) in the time instant when the specimen goes over from the s into the n state. In other words,

$$\mathcal{F}_{s0} \left(p + \frac{H_c^2}{8\pi}, T \right) = \mathcal{F}_n(p, T),$$

where $\mathcal{F}_{s0} \left(p + \frac{H_c^2}{8\pi}, T \right)$ is the specific thermodynamic potential in the absence of the magnetic field, i.e. at the pressure $p + \frac{H_c^2}{8\pi}$.

The latter Eq. permits the following expansion by topological charges $k \in \mathbf{Z}$:

$$\sum_k \mathcal{F}_{s0k} \left(p + \frac{H_c^2}{8\pi}, T \right) = \sum_k \mathcal{F}_{n,k}(p, T)$$

because the natural additivity of the specific thermodynamic potential $\mathcal{F}$. Such "topological" expansion reflects correctly the both "discrete" and "continuous" geometries proper, respectively, to the s and n states.

At $T \neq 0$, as it can be seen easily from the latter formulas, the latent heat Q accumulated course the first-order phase transition in a Type II superconductor is liberated at a temperature $T_b < T_c$ and the superconductor returns back in the Curie point T_c [29]. Herewith the specimen becomes completely superconducting.

Thus these formulas outline the definite (perhaps, initial) approach to the high-temperature superconductivity, including the account of topological properties of that (again Abelian) model. These properties, resemble, probably, those of the Ginzburg-Landau-Abrikosov model [30, 49].

The same is correct and for the NO model [27] (if consider it in the $T \rightarrow 0$ limit).

In the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac, the question about the dynamics of the first and second-order phase transitions occurring therein is, probably, more complicate than in the liquid helium II (at rest) theory.

This is associated with the discrete geometry of the (3.3) type us assumed for the initial, $SU(2)$, and residual, $U(1)$, gauge groups in the former model.

On the other hand, again "freezing" the supercooling situation at low temperatures $T \rightarrow 0$ can be observed in the Minkowskian Higgs model with vacuum BPS monopoles quantized by Dirac.

This "frozen supercooling situation" comes to coexisting superfluid potential motions and collective solid rotations inside the BPS monopole vacuum suffered the Dirac fundamental quantization [4].

Herewith collective solid rotations inside this BPS monopole vacuum proceed without "friction forces" at $T = 0$.

In this is the essence of the Josephson effect [3, 17]: at $T = 0$, any "quantum train" cannot stop (until $\theta \neq 0$) moving along closed trajectories.

Generally speaking, the task to trace in detail the dynamics of "combined" second and first-order phase transitions in gauge models is very difficult. It claims for its solving specific numerical methods (perhaps, similar to those were applied in Ref. [58] for tracing highly complicate dynamics of the first-order phase transition in GUT).

References

- [1] L. D. Lantsman, Dirac Fundamental Quantization of Gauge Theories is the Natural Way of Reference Frames in Modern Physics, submitted in Int. J. Mod. Phys. A, [arXiv:hep-th/0604004].
- [2] L. D. Lantsman, Superfluidity of Minkowskian Higgs Vacuum with BPS Monopoles Quantized by Dirac May Be Described as Cauchy Problem to Gribov Ambiguity Equation., [arXiv:hep-th/0607079].
- [3] L. D. Lantsman, Nontrivial Topological Dynamics in Minkowskian Higgs Model Quantized by Dirac., [arXiv:hep-th/0610217].
- [4] P. A. M. Dirac, Proc. Roy. Soc. A 114 (1927) 243; Can. J. Phys. 33 (1955) 650.

- [5] D. Blaschke, V. N. Pervushin, G. Röpke, Topological Gauge Invariant Variables in QCD, Proc. of Workshop: Physical Variables in Gauge Theories, JINR, Dubna, Russia, 21-24 Sept. 1999, MPG-VT-UR 191/99, [arXiv:hep-th/9909133].
- [6] L. D. Lantsman, Superfluid Properties of BPS Monopoles, [arXiv:hep-th/0605074].
- [7] L. D. Lantsman, V. N. Pervushin, The Higgs Field as The Cheshire Cat and his Yang-Mills "Smiles", Proc. of 6th International Baldin Seminar on High Energy Physics Problems (ISHEPP), Dubna, Russia, 10-15 June 2002, [arXiv:hep-th/0205252];
L. D. Lantsman, Minkowskian Yang-Mills Vacuum, [arXiv:math-ph/0411080].
- [8] L. D. Lantsman, V. N. Pervushin, Yad. Fiz. 66 (2003) 1416 [Physics of Atomic Nuclei 66 (2003) 1384], JINR P2-2002-119, [arXiv:hep-th/0407195].
- [9] A. S. Schwarz, Kvantovaja Teorija Polja i Topologija, 1st edition (Nauka, Moscow, 1989) [A. S. Schwartz, Quantum Field Theory and Topology (Springer, 1993)]; Nucl. Phys. B 208 (1982) 141.
- [10] M. K. Prasad, C. M. Sommerfeld, Phys. Rev. Lett. 35 (1975) 760;
E. B. Bogomol'nyi, Yad. Fiz. 24 (1976) 449.
- [11] R. Akhoury, Ju- Hw. Jung, A. S. Goldhaber, Phys. Rev. 21 (1980) 454.
- [12] V. N. Pervushin, Dirac Variables in Gauge Theories, Lecture Notes in DAAD Summerschool on Dense Matter in Particle and Astrophysics, JINR, Dubna, Russia, 20-31 August 2001, Phys. Part. Nucl. 34 (2003) 348 [Fiz. Elem. Chast. Atom. Yadra 34 (2003) 679], [arXiv:hep-th/0109218].
- [13] L. D. Landau, E. M. Lifschitz, Lehrbuch der Theoretischen Physik (Statistische Physik, Band 5, teil 2), in German, 1st edition by H. Eserig and P. Ziesche (Akademie-Verlag, Berlin, 1980).
- [14] L. D. Landau, JETP 11 (1941) 592; DAN USSR 61 (1948) 253.
- [15] I. M. Khalatnikov, Teorija Sverxtekychesti, 1st edition (Nauka, Moscow, 1971).
- [16] V. N. Pervushin, Teor. Mat. Fiz. 45 (1980) 394 [Theor. Math. Phys. 45 (1981) 1100].
- [17] V. N. Pervushin, Riv. Nuovo Cim. 8, N 10 (1985) 1.
- [18] L. D. Faddeev, Proc. of 4th Int. Symp. on Nonlocal Quantum Field Theory, Dubna, USSR, 1976, JINR D1-9768, p. 267.
- [19] G. 't Hooft, Nucl. Phys. B 138 (1978) 1.

- [20] D. Blaschke, V. N. Pervushin, G. Röpke, Proc. of the Int. Seminar Physical variables in Gauge Theories, Dubna, Russia, 21-24 Sept. 1999, edited by A. M. Khvedelidze, M. Lavelle, D. McMullan and V. Pervushin (E2-2000-172, Dubna, 2000), p. 49, [arXiv:hep-th/0006249].
- [21] P. A. M. Dirac, Proc. Roy. Soc. A 133 (1931) 69.
- [22] F. Bruckmann, G. 't Hooft, Phys. Rep. 142 (1986) 357; [arXiv:hep-th/0010225].
- [23] P. I. Azimov, V. N. Pervushin, Teor. Mat. Fiz. 67 (1986) 349 [Theor. Math. Phys. 67 (1987) 546].
- [24] T. P. Cheng, L.- F. Li, Gauge Theory of Elementary Particle Physics, 3rd edition (Clarendon Press, Oxford, 1988).
- [25] L. Del Debbio, M. Faber, J. Greensite, S. Olejnik, Phys. Rev. D 55 (1997) 2298, [arXiv:hep-lat/9610005].
- [26] M. A. L. Capri, D. Dudal, J. A. Gracey, V. E. R. Lemes, R. F. Sobreiro, S. P. Sorella, R. Thibes, H. Verschelde, The Infrared Behaviour of the Gluon and Ghost Propagators in SU(2) Yang-Mills Theory in the Maximal Abelian Gauge, Talk given by S.P. Sorella at the "I Latin American Workshop on High Energy Phenomenology (I LAWHEP)", 1-3 December 2005, Porto Alegre, Rio Grande Do Sul, Brasil, Instituto de Fisica, UFRGS, [arXiv:hep-th/0603167].
- [27] H. B. Nielsen, P. Olesen, Nucl. Phys. B 61 (1973) 45.
- [28] A. D. Linde, Elementary Particle Physics and Inflationary Cosmology, 1st edition (Nauka, Moscow, 1990), [arXiv: hep-th/0503203].
- [29] P. Coles, F. Lucchin, Cosmology, the Origin and Evolution of Cosmoc Structure, 2nd edition (Baffins Lane: John Wiley and Sons, LTD, 2002).
- [30] V. L. Ginzburg, L. D. Landau, JETP 20 (1950) 1064.
- [31] F. Lenz, Topological Concepts in Gauge Theories, Lectures given at the Autumn School "Topology and Geometry in Physics", of the Graduiertenkolleg "Physical systems with many degrees of freedom", Germany, University of Heidelberg, Rot an der Rot, 24-28 September 2001, FAU-TP3-04/3, [arXiv:hep-th/0403286].
- [32] A. Kronfeld, M. Laursen, G. Schrierholz, U.- J. Wiese, Phys. Lett. B 198 (1987) 516;
T. Suzuki, I. Yotsuyanagi, Phys. Rev. D 42 (1990) 4257;
S. Hioki et al., Phys. Lett. B 272 (1991) 326.
- [33] G. 't Hooft, Nucl. Phys. B 79 (1974) 276.

- [34] A. M. Polyakov, Pisma JETP 20 (1974) 247 [Sov. Phys. JETP Lett. 20 (1974) 194];
Sov. Phys. JETP Lett. 41 (1975) 988.
- [35] L. D. Faddeev, V. N. Popov, Phys. Lett. B 25 (1967) 29.
- [36] L. H. Ryder, Quantum Field Theory, 1st edition (Cambridge University Press, Cambridge, 1984).
- [37] L. M. Krauss, F. Wilczek, Phys. Rev. Lett. 62 (1989) 1221;
M.G. Alford, J. M. Russell, F. Wilczek, Nucl.Phys. B 328 (1989) 140.
- [38] M. A. Markov, J. Phys. (USSR) **2**, 453 (1940);
H. Yukawa, Phys. Rev. **77**, 219 (1950).
- [39] L. D. Landau, E. M. Lifschitz, Theoretical Physics, v. 2. The Field Theory, edited
by L. P. Pitaevskii, 7th edition (Nauka, Moscow, 1988).
- [40] R. Engelking, General Topology, 2nd edition (Państwowe Wydawnictwo Naukowe,
Warszawa, 1977).
- [41] D. R. Nelson, Defects in Superfluids, Superconductors and Membranes, Lectures
presented at the 1994 Les Houches Summer School "Fluctuating Geometries in
Statistical Mechanics and Field Theory.", [arXiv:cond-mat/9502114].
- [42] N. N. Bogoliubov, J. Phys. 9 (1947) 23;
N. N. Bogoliubov, V. V. Tolmachev, D. V. Shirkov, Novij Metod v Teorii Sverch-
provodimosti, 1st edition (Izd-vo AN SSSR, Moscow, 1958: p. p. 5-9).
- [43] V. G. Levich, Yu. A. Vdovin, V. A. Mjamlin, Kurs Teoreticheskoy Fiziki, v. 2, 2nd
edition (Nauka, Moscow, 1971).
- [44] V. N. Pervushin, V. I. Smirichinski, J. Phys. A 32 (1999) 6191, JINR-E2-99-16,
[arXiv:hep-th/9902013].
- [45] M. Bucher, Hoi-Kwong Lo, J. Preskill, Nucl. Phys. B 386 (1992) 3,
[arXiv:hep-th/9112039].
- [46] V. G. Levich, Yu. A. Vdovin, V. A. Mjamlin, Kurs Teoreticheskoy Fiziki, v. 1, 2nd
edition (Nauka, Moscow, 1969).
- [47] V. S. Vladimirov, Yravneniya Matematicheskoy Fiziki, 5th edition (Nauka, Moscow,
1988).
- [48] A. D. Linde, Phys. Lett B 96 (1980) 289.
- [49] A. A. Abrikosov, JETP 32 (1957) 1442.

- [50] L. D. Landau, E. M. Lifshitz, Theoretical Physics, v. 4. Quantum Electrodynamics (V. B. Berestetskii, E. M. Lifshitz, L. P. Pitaevskii), edited by L. P. Pitaevskii, 3rd edition (Nauka, Moscow, 1989).
- [51] A. I. Achiesser, V. B. Berestetskii, Quantum Electrodynamics, 3rd edition (Nauka, Moscow, 1969).
- [52] Y. Aharonov, D. Bohm, Phys. Rev. 115 (1959) 484.
- [53] L.D. Landau, E. M. Lifshitz, Elektrodinamika sploshnix sred (in Russian), edited by L. P. Pitaevskii, 2nd edition (Nauka, Moscow, 1982).
- [54] Yu. L. Kalinovskiy, W. Kallies, V. N. Pervushin, N. A. Sarikov, Fortschr. Phys. 38 (1990) 333.
- [55] A. A. Bogolubskaya, Yu. L. Kalinovskiy, W. Kallies, V. N. Pervushin, Acta Phys. Polonica 21 (1990) 139.
- [56] T. T. Wu, C. N. Yang, Phys. Rev. D 12 (1975) 3845.
- [57] A. A. Bykov, I. M. Dremin, A. V. Leonidov, Usp. Fiz. Nauk 143 (1984) 3 [Sov. Phys. Usp. 27 (1984) 321].
- [58] V. A. Kuzmin, M. E. Shaposhnikov, I. I. Tkachev, Z. Phys. C 12 (1982) 83.